

SUPPLEMENTARY MATERIAL

Review of the Impact of Weight Loss and Body Mass Index in Clinical Trials of Nintedanib in Interstitial Lung Disease

Michael Kreuter,^{1*} R Matthew Kottmann,² Fengming Luo,³ Carl Coeck,⁴ Madhu Kanakapura,⁵
Christina Schlecker,⁵ Ivana Ritter,⁵ Tomohiro Handa⁶

¹ Mainz Center for Pulmonary Medicine, Department of Pneumology, Mainz University Medical Center and Marienhaus Clinic Mainz, Mainz, Germany.

² Division of Pulmonary Disease and Critical Care Medicine, University of Rochester Medical Center, Rochester, NY, USA.

³ Department of Respiratory and Critical Care Medicine, West China Hospital, Sichuan University, Chengdu, China.a

⁴ Boehringer Ingelheim SComm, Brussels, Belgium.

⁵ Boehringer Ingelheim International GmbH, Ingelheim am Rhein, Germany.

⁶ Department of Advanced Medicine for Respiratory Failure, Graduate School of Medicine, Kyoto University, Kyoto, Japan.

*Corresponding author: Prof. Dr. Michael Kreuter, M.D. Mainz Center for Pulmonary Medicine, Department of Pneumology, Mainz University Medical Center and Marienhaus Clinic Mainz, Mainz, Germany. Email: kreuterm@uni-mainz.de

List of investigators for INPULSIS¹⁶

Australia: T. Corte, Royal Prince Alfred Hospital, Camperdown, New South Wales; H. Davies, Repatriation General Hospital, Daw Park, South Australia; I. Glaspole, Alfred Hospital, Melbourne, Victoria; J. Mulder, Frankston Hospital, Frankston, Victoria; E. Veitch, Concord Repatriation General Hospital, Concord, New South Wales. **Belgium:** P. De Vuyst, Erasme University Hospital, Brussels; G. Liistro, Cliniques Universitaires Saint-Luc, Brussels; Y. Sibille, Cliniques Universitaires UCL de Mont-Godinne, Yvoir; W. Vincken, Universitair Ziekenhuis Brussel, Brussels; W. Wuyts, Universitair Ziekenhuis Leuven Gasthuisberg Campus, Leuven. **Canada:** C. Fell, Peter Lougheed Centre, Calgary, Alberta; P. Hernandez, Queen Elizabeth II Health Sciences Centre, Halifax, Nova Scotia; M. Kolb, St. Joseph's Healthcare, Hamilton, Ontario. **Chile:** A. Undurraga, Instituto Nacional de Torax, Santiago. **China:** C. Bai, Zhongshan Hospital Fudan University, Shanghai; P. Chen, The General Hospital of Shenyang Military Command, Shenyang; Z. Gao, Peking University People's Hospital, Beijing; J. Kang, The First Hospital of Chinese Medical University, Shenyang; H. Li, Shanghai Pulmonary Hospital, Shanghai; Z. Li, Xijing Hospital, 4th Military Medical University, Xi'an; H. Wan, Shanghai Ruijin Hospital, Shanghai; H. Wang, Beijing Friendship Hospital, Beijing; F. Wen, West China Hospital Sichuan University, Chengdu; Q. Xiao, Xiangya Hospital, Central South University, Changsha; Z. Xu, Peking Union Medical College Hospital, Beijing; W. Zhang, The First Affiliated Hospital of Nanchang University, Nanchang; X. Zheng, General Hospital of Ningxia Medical University, Yinchuan; H. Zhu, Shanghai Huadong Hospital, Shanghai. **Czech Republic:** N. Pauk, University Hospital Na Bulovce, Praha; P. Reiterer, Masaryk Hospital, Usti nad Labem; M. Vasakova, Thomayer's Hospital, Clinic of Pneumology, Praha. **Finland:** U. Hodgson, HYKS Keuhkosairauksien tutkimusyksikkö, Helsinki. **France:** A. Bourdin, Hôpital A. Villeneuve, Pneumologie, Montpellier; J. Cadranel, Hôpital Tenon, Pneumologie, Paris; P. Camus, Hôpital du Bocage, Pneumologie, Dijon; P. Chanez, Hôpital Nord, Pneumologie, Marseille; V. Cottin, Hôpital Louis Pradel, Pneumologie, Lyon; B. Crestani, Hôpital Bichat, Pneumologie, Paris; D. Israel-Biet, Hôpital Européen Georges-Pompidou, Pneumologie, Paris; S. Jouneau, Hôpital Pontchaillou, Pneumologie, Rennes; F. Lebargy, Hôpital Maison Blanche, Pneumologie, Reims; C. H. Marquette, Hôpital Pasteur, Pneumologie, Nice; G. Prévot, Hôpital Larrey, Pneumologie, Toulouse; D. Valeyre, Hôpital Avicenne, Pneumologie, Bobigny; B. Wallaert, Hôpital Calmette, Pneumologie, Lille. **Germany:** R. Bonnet, Zentralklinik Bad Berka GmbH, Bad Berka; U. Costabel, Ruhrlandklinik, Essen; S. Gläser, Ernst-Moritz-Arndt-Universität, Greifswald; C. Grohé, Evangelische Lungenklinik Berlin; A. Guenther, Universitätsklinikum Gießen, Gießen; P. Hammerl, Fachklinik für Lungenerkrankungen, Immenhausen; G. Höffken, Uniklinik Carl Gustav Carus, Dresden; C. Karagiannidis, Kliniken der Stadt Koeln, Koeln; J. Kirschner, CIMS Studienzentrum Bamberg UG, Bamberg; A. Kirsten, LungenClinic, Grosshansdorf; S. Korn, Johannes-Gutenberg-Universitätsklinik, Mainz; M. Kreuter, Universitätsklinikum Heidelberg, Heidelberg; J. Müller-Quernheim, Klinik der Albert-Ludwig-Universität, Freiburg; C. Neurohr, Klinikum Großhadern der L.M.-Universität, München; M. Pfeifer, Krankenhaus, Donaustauf; N. Schönfeld, HELIOS-Kliniken Emil von Behring, Berlin; R. Wiewrodt, Universitätsklinikum Münster, Medizinischen Poliklinik D, Münster. **Greece:** K. Antoniou, University Hospital of Heraklion, University Pulmonology CI, Heraklion, Crete; Z. Daniil, University Hospital of Larissa, Pulmonology Clinic, Larissa; F. Diamantea, General Hospital of Athens "Sismanoglio", 3rd Pneumonology Department, Athens; N. Koulouris, Athens Hospital of Chest Diseases "Sotiria", Athens; G. Mathioudakis, General Hospital of Nikaia, Pulmonology Clinic, Nikaia. **India:** A. G. Ghosal, National Allergy Asthma Bronchitis Institute, Kolkata; S. Kadappa Shivappa, Chest and Maternity Centre, Bangalore; M. M. Kawedia, Jehangir Hospital, Pune; P. Khatavkar, King Edward Memorial Hospital, Mumbai; A. Kumar, Asthma Bhawan, Jaipur; P. Mehta, Mehta Hospital and Cardiopulmonary

Care Centre, Ahmedabad; V. Singh, SMS Medical College and Hospital, Jaipur; K. Srikanth, PSG Hospitals, Coimbatore; H. Thakker, Qualysure Clinical Research Solution, Ahmedabad; Z. Udwadia, P.D. Hinduja National Hospital and Medical Research Center, Mumbai.

Ireland: J. Egan, Mater Misericordiae University Hospital, Department of Respiratory Medicine, Dublin. **Israel:** G. Fink, Kaplan Medical Center, Pulmonary Institute, Rehovot; M. R. Kramer, Rabin Medical Center, Petah Tiqwa; M. Yigla, Rambam Medical Center, Haifa.

Italy: C. Agostini, Università degli Studi, Dipartimento di Medicina Clinica e Sperimentale, Padova; F. De Benedetto, P.O. Clinicizzato SS. Annunziata, U.O. di Pneumologia, Chieti; S. Harari, Ospedale S. Giuseppe Fatebenefratelli, Pneumologia, Milano; F. Luppi, Università Modena e Reggio Emilia, Malattie dell'Apparato Respiratorio, Modena; P. L. Paggiaro and L. M. Tavanti, Azienda Ospedaliera Universitaria Pisana, Ospedale Cisanello, Fisiop. Respiratoria, Pisa; A. Pesci, Nuovo Ospedale "S. Gerardo", U.O. Pneumologia, Monza; V. Poletti, Ospedale Morgagni-Pierantoni, Dipartimento Toracico U.O. Pneumologia, Forlì; P. Rottoli, Presidio Ospedale Le Scotte, Clinica Pneumologica, Siena; C. Saltini, Policlinico Università Tor Vergata, Malattie Respiratorie, Roma; A. Sanduzzi Zamparelli, Università Federico II – II Clinica Pneumologica, Napoli; C. Vancheri, A.O.U. Policlinico Dipartimento di Medicina Interna e Special. Sez. Mal. Resp., Catania. **Japan:** M. Bando, Jichi Medical University Hospital, Pulmonary Medicine, Tochigi; Y. Hasegawa, Nagoya University Hospital, Respiratory Medicine, Aichi; K. Hashimoto, Tottori University Hospital, Respiratory Medicine, Tottori; S. Homma, Toho University Omori Medical Center, Respiratory Medicine, Tokyo; N. Inase, Tokyo Medical and Dental University, Pulmonary Medicine, Tokyo; Y. Inoue and T. Arai, National Hospital Organization Kinki-Chuo Chest Medical Center, Department of Internal Medicine, Osaka; S. Izumi, National Center for Global Health and Medicine, Respiratory Medicine, Tokyo; T. Kawamura, Himeji Medical Center, Respiratory Tract Medicine, Hyogo; K. Kishi, Toranomon Hospital, Department of Respiratory Medicine, Respiratory Center, Tokyo; Y. Kondo, Tosei General Hospital, Department of Allergy and Respiratory Medicine, Aichi; K. Kuwano, The Jikei University Hospital, Division of Respiratory Disease, Department of Internal Medicine, Tokyo; Y. Miura, Nippon Medical School Hospital, Respiratory Medicine, Tokyo; Y. Nishioka, Tokushima University Hospital, Respiratory Medicine and Rheumatology, Tokushima; O. Nishiyama, Kinki University Hospital, Respiratory Medicine and Allergy, Osaka; T. Ogura, Kanagawa Cardiovascular and Respiratory Center, Respiratory Medicine, Kanagawa; S. Ohkouchi, Tohoku University Hospital, Department of Respiratory Medicine, Miyagi; T. Saito, Ibaraki Higashi National Hospital, Ibaraki Department of Respiratory Medicine, Naka; Y. Setoguchi, Tokyo Medical University Hospital, Department of Respiratory Medicine, Respiratory Center, Tokyo; J. Shindoh, Ogaki Municipal Hospital, Respiratory Med, Gifu; Y. Taguchi, Tenri Hospital, Department of Respiratory Medicine, Nara; M. Tanakadate, Toranomon Hospital Kajigaya, Pulmonary and Critical Care Medicine, Kanagawa; K. Tomii, Kobe City Medical Center General Hospital, Respiratory Medicine, Hyogo; Y. Sugita, Saitama Cardiovascular and Respiratory Center, Respiratory Medicine, Saitama; T. Yamaguchi, JR Tokyo General Hospital, Respiratory Medicine, Tokyo; K. Yoshimori, Fukujūji Hospital, Respiratory Medicine, Tokyo. **Republic of Korea:** S. H. Jeong, Gachon University Gil Hospital, Incheon; D. S. Kim, Asan Medical Center, Seoul; Y. W. Kim, Seoul National University Hospital, Seoul; C. S. Park, Soon Chun Hyang University Hospital Bucheon, Bucheon; J. S. Song, The Catholic University of Korea, Yeouido St. Mary's Hospital, Seoul; S. T. Uh, Soon Chun Hyang University Hospital Seoul, Seoul. **Mexico:** M. Selman, Instituto Nacional de Enfermedades Respiratorias, Ciudad de Mexico. **Netherlands:** P. Bresser, Onze Lieve Vrouwe Gasthuis, Amsterdam; J. C. Grutters, St. Antonius Ziekenhuis, Nieuwegein, Utrecht; M. S. Wijsenbeek, Erasmus Medisch Centrum, Rotterdam. **Portugal:** A. Arrobas, Centro Hospitalar e Universitário de Coimbra, EPE, Centro Hospitalar Coimbra, Pneumology Department, Coimbra; J. Cardoso, Centro Hospitalar Lisboa Central, EPE, Hospital de Santa Marta,

Pneumology Department, Lisboa; R. Costa, Hospital Fernando Fonseca, EPE, Pneumology Department, Amadora; A. Morais, Centro Hospitalar São João, EPE, Pneumology Department, Porto; S. Neves, Centro Hospitalar de Vila Nova Gaia/Espinho, Pneumology Department, Vila Nova de Gaia; M. Serrado, Centro Hospitalar Lisboa Norte, EPE, Hospital Pulido Valente, Pneumology II Department, Lisboa. **Russian Federation:** M. Ilkovick, Scientific Research Institute of Pulmonology, Saint Petersburg; A. Vizel, GOU VPO Kazan State Medical University of Roszdrav, Kazan. **Spain:** I. Alfageme Michavila, Hospital de Valme, Sevilla; J. Ancochea, Hospital Universitario de la Princesa, Madrid; D. Castillo Villegas, Hospital de la Santa Cruz y San Pablo, Barcelona; M. Molina-Molina, Ciutat Sanitaria Universitaria Bellvitge, Barcelona; F. Morell, Hospital Vall d'Hebron, Barcelona; A. Xaubet, Hospital Clinic i Provincial de Barcelona, Neumología, Barcelona.

Türkiye: S. Aktogu Ozkan, Dr. Suat Seren Gogus Hastaliklari ve Gogus Cerrahisi Egit ve Arastirma Hastanesi, Konak/İzmir; O. Kayacan, Ankara Universitesi Tip Fakültesi Göğüs Hastaliklari, Ankara; G. Ongen, Istanbul Universitesi Cerrahpasa Tip Fakultesi Gogus Hastaliklari, Istanbul; N. Mogulkoc, Ege Universitesi Tip Fakultesi Göğüs Hastaliklari ABD, İzmir; E. Tuncay, Yedikule Gogus Hastaliklari ve Gogus Cer Egitim ve Arastirma Hastanesi, Istanbul. **United Kingdom:** P. Beirne, St James's University Hospital, Respiratory Medicine, Leeds; H. Bettinson, Churchill Hospital, Oxford Centre for Respiratory Medicine, Oxford; P. Burge, Birmingham Heartlands Hospital, Department of Respiratory Medicine, Birmingham; O. Dempsey, Aberdeen Royal Infirmary, Respiratory Department, Aberdeen; T. Maher, Royal Brompton Hospital, Interstitial Lung Disease Unit, London; A. Millar, Southmead Hospital, North Bristol Lung Centre, Bristol; L. Spencer, University Hospital Aintree, Aintree Chest Centre, Liverpool; D. Thickett, Queen Elizabeth Hospital, Lung Investigation Unit, Birmingham.

United States of America: J. Alvarez, Loess Hill Research Center, Council Bluffs, Iowa; C. Andrews, Diagnostics Research Group, San Antonio, Texas; O. Bajwa, Allegheny General Hospital, Allegheny-Singer Research Institute, Pittsburgh, Pennsylvania; A. Baker, Lynchburg Pulmonary Associates, Lynchburg, Tennessee; R. Baughman, University of Cincinnati Medical Center, Cincinnati, Ohio; J. Belperio, David Geffen School of Medicine at UCLA, Los Angeles, California; J. Bradley, Consultants in Pulmonary Medicine, Olathe, Kansas; H. Collard, University of California at San Francisco, San Francisco, California; F. Cordova, Temple Lung Center Ambulatory Care Center, Philadelphia, Pennsylvania; C. Daniels, Mayo Clinic College of Medicine, Rochester, New York; J. de Andrade, University of Alabama, Division of Pulmonary, Birmingham, Alabama; K. Dushay, Rhode Island Hospital, Providence, Rhode Island; R. Enelow, Dartmouth Hitchcock Medical Center, Lebanon, Pennsylvania; N. Ettinger, Cardiopulmonary Associates, Chesterfield, Missouri; K. Gibson, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania; M. Gotfried, Pulmonary Associates Clinical Trials, Phoenix, Arizona; A. Hajari Case, Georgia Clinical Research, Austell, Georgia; D. Hotchkin, Oregon Clinic, PC, Portland, Oregon; J. T. Huggins, Medical University of South Carolina, Charleston, South Carolina; M. Kaye, Minnesota Lung Center, Ltd., Minneapolis, Minnesota; C. Kershaw, University of Texas Southwestern Medical Center, Dallas, Texas; S. Kureishy, Metroplex Pulmonary and Sleep Centre, McKinney, Texas; L. Lancaster, Vanderbilt University Medical Center, Nashville, Tennessee; D. Lederer, New York Presbyterian Hospital, New York, New York; Y. Mageto, Vermont Lung Center, Colchester, Vermont; J. Masson, Pulmonary and Critical Care Associates, Albany, New York; K. Meyer, University of Wisconsin Hospital and Clinics, Madison, Wisconsin; P. Mohabir, Stanford Medical Center Pulmonary and Critical Care Clinic, Stanford, California; L. Morrison, Duke University Medical Center, Durham, North Carolina; S. Nathan, Inova Fairfax Hospital, Falls Church, Virginia; I. Noth, University of Chicago, Chicago, Illinois; D. Oelberg, Western Connecticut Medical Group P.C., Danbury, Connecticut; F. Rahaghi, Cleveland Clinic Florida, Weston, Florida; D. Riley, University of Medicine and Dentistry of NJ, New Brunswick, New Jersey; A. Rizzo, Lung Health and Sleep

Enhancement Center, LLC, Newark, New Jersey; M. Rossman, Hospital of the University of Pennsylvania, Philadelphia, Pennsylvania; J. Ruzi, Arizona Pulmonary Specialists, Scottsdale, Arizona; P. Sachs, Pulmonary Associates of Stamford, Stamford, Connecticut; T. Schaumberg, Oregon Clinic, PC, Portland, Oregon; M. Scholand, Lung Health Research Center, Salt Lake City, Utah; C. Schroeder, Sansum Clinic, Santa Barbara, California; F. Seifer, FDC Seifer PLC Pulmonary, Shelbyville, Illinois; J. Shea, DCOL Center for Clinical Research, Longview, Texas; D. Sinkowitz, UCLA David Geffen School of Medicine, Torrance, California; J. Tabak, Miami Research Associates, South Miami, Florida; J. Taylor, MultiCare Pulmonary Specialists, Tacoma, Washington; J. Thompson, Baptist Pulmonary and Critical Care Associates, Lexington, Kentucky; C. Thurm, Jamaica Hospital Medical Center, Department of Medicine, Jamaica, New York; J. Tita, ID Clinical Research, Ltd., Toledo, Ohio; M. Wencel, Via Christi Clinic, PA, Wichita, Kansas; J. Westerman, Jasper Summit Research, LLC, Jasper, Alabama.

List of investigators for INBUILD³¹

Argentina: S. Quadrelli, Fundación Sanatorio Güemes, Buenos Aires; M. Otaola, Centro de Investigaciones Metabólicas (CINME), Buenos Aires; M. A. Bergna, CEMER-Centro Medico De Enfermedades Respiratorias, Florida; P. Elias, INSARES, Mendoza; G. Arce, Instituto Médico de la Fundación Estudios Clínicos, Rosario; A. Cazaux, Centro Dr. Lazaro Langer S.R.L, Alberdi Sur; **Belgium:** W. Wuyts, UZ Leuven, Leuven; J. Guiot, CHU Sart Tilman, Angleur; B. Bondue, Erasme University Hospital, Bruxelles; C. Dahlqvist, Cliniques universitaires UCL de Mont-Godinne, Yvoir; **Canada:** L. Homik, Concordia Hospital Respiratory Research, Winnipeg; S. Shapera, Toronto General Hospital, Toronto; A. Cantin, CHUS-Fleurimont Centre Recherche Clinique, Sherbrooke; M. Kolb, St. Joseph's Healthcare Firestone Institute for Respiratory Health, Hamilton; **Chile:** M. Salinas Fénero, Instituto Nacional del Tórax Departamento de Neumonología, Providencia, Santiago; R. Maturana Rozas, Hospital Clínico Regional de Concepción Dr. Guillermo Grant Benavente, Concepción; A. Silva Orellana, Centro de Investigación del Maule, Talca; **China:** Z. Xu, Peking Union Medical College Hospital, Beijing; Q. Luo, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou; J. Kang, The First Hospital of China Medical University, Shenyang; H. Cai, Nanjing Drum Tower Hospital Neurology, Nanjing; **France:** S. Marchand-Adam, Hôpital Bretonneau – CHRU de Tours, Tours; E. Bergot, Hôpital Côte de Nacre – CHU de Caen, Caen; A. Gamez-Dubuis, Hôpital Arnaud de Villeneuve – CHRU de Montpellier, Montpellier; F. Riviere, Hôpital d'Instruction des Armées Percy, Clamart; R. Kessler, Hôpital Civil – Nouvel Hôpital Civil – CHU de Strasbourg, Strasbourg; H. Nunes, Hôpital Avicenne, Bobigny; C. Marquette, Hôpital Pasteur – CHU de Nice, Nice; L. Wemeau, Hôpital Calmette – CHRU de Lille, Lille; S. Jouneau, Hôpital Pontchaillou – CHU de Rennes, Rennes; F. Lebargy, Hôpital Maison Blanche – CHU de Reims, Reims; B. Crestani, Hôpital Bichat – AP-HP, Paris; V. Cottin, Hôpital Louis Pradel – Groupement Hospitalier Est – CHU de Lyon HCL, Bron; M. Reynaud-Gaubert, Hôpital Nord – AP-HM, Marseille; **Germany:** S. Blaas, Klinik Donaustauf Zentrum für Pneumologie, Donaustauf; F. Bonella, Ruhrlandklinik Westdeutsches Lungenzentrum, Essen; W. Randerath, Krankenhaus Bethanien GmbH, Solingen; J. Hetzel, Universitätsklinikum Tuebingen, Tuebingen; D. Koschel, Fachkrankenhaus Coswig GmbH, Coswig; M. Kreuter, Thoraxklinik-Heidelberg gGmbH am Universitätsklinikum Heidelberg, Heidelberg; A. Prasse, Medizinische Hochschule Hannover, Hannover; D. Skowasch, Universitätsklinikum Bonn AöR, Bonn; S. Stieglitz, Petrus Krankenhaus Wuppertal, Wuppertal; **Italy:** R. Refini, Azienda Ospedaliera Universitaria Senese, Siena; S. Cerri, Clinica Malattie Apparato Respiratorio Policlinico di Modena, Modena; A. Pesci, Azienda Ospedaliera San Gerardo di Monza U.O. Clinica Pneumologica, Monza; S. Tomassetti, AUSL di Forlì U.O., Forlì; C. Vancheri, A.O.U.

Policlinico Vittorio Emanuele U.O.P.I. interstiziopatie e malattie rare del polmone, Catania; F. Varone, Policlinico Universitario A. Gemelli U.O.C. Pneumologia, Rome;

Japan: N. Sakamoto, Nagasaki University Hospital, Nagasaki; S. Abe and H. Hayashi, Nippon Medical School Hospital, Tokyo, Bunkyo-ku; T. Saito, Ibarakihigashi National Hospital, Ibaraki, Naka-gun; T. Suda, Hamamatsu University Hospital, Shizuoka, Hamamatsu; H. Kitamura, Kanagawa Cardiovascular and Respiratory Center, Kanagawa, Yokohama; M. Okamoto, Kurume University Hospital, Fukuoka, Kurume; Y. Kondoh, Tosei General Hospital, Aichi, Seto; S. Makino and T. Takeuchi, Osaka Medical College Hospital, Osaka, Takatsuki; Y. Yamada and C. Kono, JR Tokyo General Hospital, Tokyo, Shibuya-ku; Y. Inoue, National Hospital Organization Kinki-Chuo Chest Medical Center, Osaka, Sakai; H. Sugiura, Tohoku University Hospital, Miyagi, Sendai; K. Kishi and H. Takaya, Toranomon Hospital, Tokyo, Minato-ku; H. Yamauchi, Jichi Medical University Hospital, Tochigi, Shimotsuke; K. Ichikado, Saiseikai Kumamoto Hospital, Kumamoto; K. Tomii, Kobe City Hospital Organization Kobe City Medical Center General Hospital, Hyogo, Kobe; H. Takahashi, Sapporo Medical University Hospital, Hokkaido, Sapporo; S. Izumi, National Center for Global Health and Medicine, Tokyo, Shinjuku-ku; T. Kawamura, National Hospital Organization Himeji Medical Center, Hyogo, Himeji; Y. Nishioka, Tokushima University Hospital, Tokushima; Y. Miyazaki, Tokyo Medical and Dental University, Tokyo, Bunkyo-ku;

Republic of Korea: J.W. Song, Asan Medical Center, Seoul; J.S. Park, Seoul National University Bundang Hospital, Seongnam; Y. Kim, The Catholic University of Korea, Bucheon St.Mary's Hospital, Bucheon; **Poland:** E. Jassem, University Clinical Center, Gdansk; J. Kus, National Institute of Tuberculosis and Lung Diseases Outpatient Clinic, Warsaw; W. Piotrowski, Department of Pneumology and Allergy, Medical University of Lodz, Lodz; A. Barczyk, Leszek Giec Upper-Silesian Medical Centre, Katowice; D. Ziora, Dept & Clinic of Internal Diseases/Pulmon.dept, Zabrze; **Russian Federation:** E. Bazdyrev, Res.Inst.- Compl.Iss.Cardi.Dis., Kemerovo; S. Moiseev, Sechenov First Moscow State Medical University, Moscow; S. Avdeev, Sechenov First Moscow State Medical University, Moscow; M. Ilkovich, Scientific Research Institute of Pulmonology Pavlov State Medical University Clinical Center f. Inst, St. Petersburg; V. Yakusevich, Clinical Hospital of Emergency Care n.a. N.V. Solovyev, Yaroslavl; **Spain:** C. Valenzuela, Hospital Universitario de La Princesa, Madrid; O. Acosta, Hospital Universitario de Canarias, San Cristóbal de La Laguna; M. Martínez, Hospital Universitari i Politècnic La Fe, Valencia; L. Gómez Carrera, Hospital Universitario La Paz, Madrid; M. Molina-Molina, Hospital Universitari de Bellvitge, L'Hospitalet de Llobregat; D. M. Castillo Villegas, Hospital de la Santa Creu i Sant Pau, Barcelona; M. Aburto, Hospital de Galdakao, Galdakao; J.A. Rodríguez Portal, Hospital Universitario Virgen del Rocío, Sevilla; A. Villar, Hospital Universitari Vall d'Hebron, Barcelona; A. León Jiménez, Hospital Puerta del Mar, Cádiz; J. Saulea, Hospital Son Espases, Palma de Mallorca; M. Arias, Hospital Central de Asturias, Oviedo;

United Kingdom: T. M. Maher, Royal Brompton Hospital, London; P. Beirne, St James's University Hospital, Leeds; H. Stone, Royal Stoke University Hospital, Stoke-on-Trent; B. Hope-Gill, University Hospital Llandough, Cardiff; N. Hirani, Royal Infirmary of Edinburgh, Edinburgh; N. Chaudhuri, Wythenshawe Hospital, Manchester; **United States of America:** C. Andrews, Diagnostics Research Group, San Antonio; A. Gifford, Dartmouth-Hitchcock Medical Center, Lebanon; L. Jones, University of Florida College of Medicine, Jacksonville; L. Morrison, Duke University Medical Center, Durham; D. Antin-Ozerkis, Yale University School of Medicine, New Haven; N. Bhatt, The Ohio State University Wexner Medical Center, Columbus; T. Kulkarni, University of Alabama at Birmingham Lung Health Center, Birmingham; T. Moua, Mayo Clinic, Rochester; N. Ettinger, The Lung Research Center, Chesterfield; L. Pitts, University of Kansas Medical Center, Kansas City; S. Veeraraghavan, The Emory Clinic, Atlanta; M. Padilla, Icahn School of Medicine at Mount Sinai – Mount Sinai Medical Center, New York; E. R. Fernández Pérez, National Jewish Health, Denver;

G. Giessel, Pulmonary Associates of Richmond, Inc., Richmond; M. Strek, University of Chicago Medical Center, Chicago; S. Danoff, Johns Hopkins Asthma and Allergy Center, Baltimore; J. Burk, Texas Pulmonary and Critical Care Consultants, PA, Fort Worth; M. Rossman, The Hospital of the University of Pennsylvania, Philadelphia; N. Patel, Columbia University Medical Center-New York Presbyterian Hospital, New York; E. Belloli, Pulmonary and Critical Care Medicine, Ann Arbor; D. Hotchkin, The Oregon Clinic, Portland; S. Weigt, Peter Morton Medical Building, Los Angeles; M. B. Scholand, University of Utah Health, Salt Lake City; R. Kaner, New York-Presbyterian Hospital/ Weill Cornell Medical Center, New York; B. Sigal, Southeastern Research Center, Winston-Salem; Z. Safdar, Houston Methodist Hospital, Houston; L. Tolle, Cleveland Clinic, Cleveland; R. Martinez, Pulmonary and Sleep of Tampa Bay, Brandon; M. Glassberg, University of Miami Pulmonary Research Office, Miami; R. Hallowell, Beth Israel Deaconess Medical Center, Boston; J. Golden, University of California San Francisco, San Francisco; M. Schwartz, Pulmonary and Critical Care Associates of Baltimore, Towson; E. Britt, University of Maryland Medical Center, Baltimore; L. Morrow, Creighton University School of Medicine, Omaha; Y. Mageto, Baylor University Medical Center, Dallas; K. Buch and S. Chaaban, University of Kentucky Medical Center, Lexington; H. Poonyagariyagorn, The Oregon Clinic – Pulmonary, Critical Care and Sleep Medicine – West, Portland; D. Dilling, Loyola University Medical Center, Maywood; O. Shlobin, Inova Fairfax Medical Campus, Falls Church; K. Thavarajah, Henry Ford Health System, Detroit; A. Nambiar, Medical Arts and Research Center (MARC), San Antonio; I. Rosas, Brigham and Women's Hospital, Boston; R. Bascom, Penn State Milton S. Hershey Medical Center, Hershey; J. Oldham, UC Davis Medical Center, Sacramento; S. Schmidt, Spectrum Health, Grand Rapids; J. Dematte D'Amico, Northwestern Memorial Hospital, Chicago; J. Falk, Cedars-Sinai Medical Center, Los Angeles; C. Glazer, UT Southwestern Medical Center, Dallas; G. Criner, Temple University Hospital, Oaks.

REFERENCES

16. Richeldi, L. et al. Efficacy and safety of nintedanib in idiopathic pulmonary fibrosis. *N Engl J Med.* 370, 2071–82 (2014).
31. Flaherty K. R. et al. Nintedanib in progressive fibrosing interstitial lung diseases. *N Engl J Med.* 381,1718–27 (2019).