

Rocky-shore polychaetes in southeastern Brazil: diversity, zonation, and human influence

Alexia de Lima Domingues da Cruz^{12*}. Breno Monzato Gisbert¹. Patrick Targino Braga Pessoa¹. Cinthya Simone Gomes Santos¹².

¹Laboratório de Sistemática e Ecologia de Polychaeta, Universidade Federal Fluminense, Niterói, Rio de Janeiro, Brazil

² Programa de Pós-Graduação em Biologia Marinha e Ambientes Costeiros, Universidade Federal Fluminense, Niterói, Rio de Janeiro, Brazil

* Corresponding author. E-mail address: alexiadomingues@id.uff.br

Supplementary Material

Table S1. Beta-diversity partitioning (presence–absence, Sørensen) across spatial scales in rocky-shore polychaete assemblages from Itaipu and Sacristia. Abbreviations: β SOR = Sørensen dissimilarity (total beta diversity); β SIM = turnover component; β SNE = nestedness component; Prop_turnover = β SIM/ β SOR; Prop_nestedness = β SNE/ β SOR; level = comparison type (“Between” = among units at that spatial scale; “Within” = average heterogeneity within units at that scale).

Scale	level	β SIM	β SNE	β SOR	Prop_turnover	Prop_nestedness
Itaipu x Sacristia	Between	0,5591	0,1577	0.7169	0.78	0.22
Itaipu x Sacristia	Within	0.4371	0.1894	0.6266	0.70	0.30
Itaipu - Stations	Between	0.3648	0.3074	0.6723	0.54	0.46
Itaipu - Stations	Within	0.4516	0.1422	0.5939	0.76	0.24
Itaipu - Zones	Between	0.4534	0.1978	0.6513	0.70	0.30
Itaipu - Zones	Within	0.3452	0.2737	0.6190	0.56	0.44
Sacristia - Stations	Between	0.4785	0.1389	0.6175	0.77	0.23
Sacristia - Stations	Within	0.4599	0.1554	0.6153	0.75	0.25
Sacristia - Zones	Between	0.5072	0.1329	0.6402	0.79	0.21
Sacristia - Zones	Within	0.4254	0.1626	0.5881	0.72	0.28

Table S2. Total abundance (n) of polychaete families and species at the rocky-shores of Itaipu and Sacristia (Niterói, Brazil). partitioned by station (1–2) and shore zone (*midlittoral*. *upper infralittoral*). Values are sums of individuals across all replicates within each stratum; zeros denote absence. Abbreviations: sp. = species undetermined; indet. = higher-rank identification; cf. = confer; juv. = juvenile; epit. = epitoke.

Family	Species	Itaipu				Sacristia			
		Station 1		Station 2		Station 1		Station 2	
		Mid	Infra	Mid	Infra	Mid	Infra	Mid	Infra
Arenicolidae	<i>Branchiomaldane</i>								
	sp.	0	0	0	0	0	0	0	5
	Arenicolidae indet.	0	0	0	0	0	0	0	23
Capitellidae	<i>Barantolla</i> sp.	0	0	2	2	0	0	0	0
	<i>Capitella</i> sp.1	0	0	1	0	0	0	1	0
	<i>Capitella</i> sp.2	0	0	0	0	0	0	3	0
	<i>Capitellethus</i> sp.	1	0	0	1	0	0	0	0
	Capitellidae indet.								
	juv.	1	0	2	0	0	0	0	0
	<i>Leiocapitellides</i> cf.								
	<i>analís</i>	0	1	0	0	0	0	0	0
	<i>Notomastus</i> sp.	0	0	1	0	0	0	0	0
	<i>Parheteromastides</i>								
	sp.	0	0	0	1	0	0	1	0
	<i>Pseudoleiocapitella</i>								
	cf. <i>fauveli</i>	0	0	9	4	0	0	0	0
	<i>Scyphoproctus</i> sp.	0	0	1	0	0	0	0	0
Cirratulidae	Cirratulidae indet.	2	0	0	8	7	10	2	11
Dorvilleidae	<i>Ophryotrocha</i> sp.	0	0	1	0	0	0	0	0
Eunicidae	<i>Eunice</i> sp.	0	0	1	0	0	0	2	0
	Eunicidae								
	indet.juv.	2	0	0	1	1	2	0	1
	<i>Lysidice ninetta</i>	0	0	0	1	0	0	0	0
	<i>Lysidice</i> sp.	0	0	0	0	1	0	0	0
	<i>Lysidice unicornis</i>	0	0	0	0	1	2	2	1
Lumbrineridae	<i>Lumbrineris</i> sp.	1	0	0	0	0	0	0	0
Nereididae	Nereididae indet.	0	2	101	16	3	4	19	2
Orbiniidae	Orbiniidae indet.	29	3	10	12	9	10	0	1
Sabellariidae	<i>Phragmatopoma</i>								
	sp.	0	0	0	1	1	0	0	0
Sabellidae	Sabellidae indet.	5	0	4	177	31	5	1	10
Saccocirridae	<i>Pharyngocirrus</i>								
	<i>gabriellae</i>	0	3	0	0	2	1	0	0
	<i>Saccocirrus</i>								
	<i>pussicus</i>	1	0	0	0	0	0	0	0
Serpulidae	<i>Serpula</i> sp.	2	0	1	1	0	1	0	0
Spionidae	Spionidae indet.	5	36	3	42	0	0	1	0
Syllidae	<i>Brania pusilla</i>	0	0	0	11	7	2	1	2
	<i>Erinaceusyllis</i>								
	<i>belizensis</i>	0	0	0	2	2	2	10	0
	<i>Eusyllis</i> sp.	0	0	2	0	0	0	1	0
	<i>Exogone</i>								
	<i>breviantennata</i>	0	1	7	40	8	18	0	19
	<i>Exogone simplex</i>	0	0	0	0	3	12	2	28
	<i>Lamellisyllis</i> sp.	0	1	0	0	0	0	0	0
	<i>Opisthosyllis</i>								
	<i>brunnea</i>	0	0	19	4	0	0	0	0
	<i>Parexogone</i>								
	<i>anseforbansensis</i>	0	0	0	2	0	0	0	0
	<i>Perkinsyllis</i> sp.	0	0	0	1	0	0	0	0
	Syllidae indet. epit.	0	1	0	0	2	0	1	0

	<i>Syllidae</i> indet. juv.	1	0	15	4	0	3	38	5
	<i>Syllis</i> cf. <i>gerlachi</i>	0	0	0	0	4	0	22	3
	<i>Syllis</i> cf. <i>prolifera</i>	1	0	0	0	0	0	0	0
	<i>Syllis hyllebergi</i>	0	3	0	10	3	6	13	2
	<i>Syllis magellanica</i>	0	0	0	4	12	4	3	3
	<i>Syllis</i> sp.	1	3	9	18	0	1	90	2
	<i>Trypanosyllis</i>								
	<i>auranticus</i>	6	1	4	5	1	0	6	0
	<i>Trypanosyllis</i> sp.1	0	0	2	1	3	0	6	2
	<i>Trypanosyllis</i> sp.2	0	0	0	0	0	0	27	0
	<i>Trypanosyllis zebra</i>	0	0	0	2	0	0	0	0
Terebellidae	Terebellidae indet.	0	1	14	43	0	0	0	0

Table S3. Relative abundance (%) of polychaete families and species at the rocky-shores of Itaipu and Sacristia, partitioned by station (1–2) and shore zone (midlittoral, upper infralittoral). The “All sites (%)” column is computed as $RA_i = n_i / \sum n \times 100$ across all strata. Zeros denote absence. Abbreviations: sp. = species undetermined; indet. = higher-rank identification; cf. = confer; juv. = juvenile; epit. = epitoke.

Family	Species	All sites (%)	Itaipu				Sacristia			
			Station 1		Station 2		Station 1		Station 2	
			Mid	Infra	Mid	Infra	Mid	Infra	Mid	Infra
Arenicolidae	<i>Branchiomaldane</i> sp.	0.39	0	0	0	0	0	0	0	4.17
Capitellidae	Arenicolidae indet.	1.78	0	0	0	0	0	0	0	19.17
	<i>Barantolla</i> sp.	0.31	0	0	0.96	0.48	0	0	0	0
	<i>Capitella</i> sp.1	0.15	0	0	0.48	0	0	0	0.4	0
	<i>Capitella</i> sp.2	0.23	0	0	0	0	0	0	1.19	0
	<i>Capitellethus</i> sp.	0.15	1.72	0	0	0.24	0	0	0	0
	Capitellidae indet.	0.08	1.72	0	0	0	0	0	0	0
	juv.									
	<i>Leiocapitellides</i> cf. <i>analis</i>	0.08	0	1.79	0	0	0	0	0	0
	<i>Notomastus</i> sp.	0.08	0	0	0.48	0	0	0	0	0
	<i>Parheteromastides</i> sp.	0.15	0	0	0	0.24	0	0	0.4	0
	<i>Pseudoleiocapitella</i> cf. <i>fauveli</i>	1.01	0	0	4.31	0.97	0	0	0	0
	<i>Scyphoproctus</i> sp.	0.08	0	0	0.48	0	0	0	0	0
Cirratulidae	Cirratulidae indet.	3.25	3.45	0	0.96	1.93	6.93	12.05	0.79	9.17
Dorvilleidae	<i>Ophryotrocha</i> sp.	0.08	0	0	0.48	0	0	0	0	0
Eunicidae	<i>Eunice</i> sp.	0.23	0	0	0.48	0	0	0	0.79	0
	<i>Eunicidae</i> indet.juv.	0.54	3.45	0	0	0.24	0.99	2.41	0	0.83
	<i>Lysidice ninetta</i>	0.08	0	0	0	0.24	0	0	0	0
	<i>Lysidice</i> sp.	0.08	0	0	0	0	0.99	0	0	0
	<i>Lysidice unicornis</i>	0.46	0	0	0	0	0.99	2.41	0.79	0.83
Lumbrineridae	<i>Lumbrineris</i> sp.	0.08	1.72	0	0	0	0	0	0	0
Nereididae	Nereididae indet.	11.37	0	3.57	48.33	3.86	2.97	4.82	7.54	1.67
Orbiniidae	Orbiniidae indet.	5.72	50	5.36	4.78	2.9	8.91	12.05	0	0.83
Sabellariidae	<i>Phragmatopoma</i> sp.	0.15	0	0	0	0.24	0.99	0	0	0
	Sabellidae									
Saccocirridae	Sabellidae indet.	18.02	8.62	0	1.91	42.75	30.69	6.02	0.4	8.33
	<i>Pharyngocirrus gabriellae</i>	0.46	0	5.36	0	0	1.98	1.2	0	0
	<i>Saccocirrus pussicus</i>	0.08	1.72	0	0	0	0	0	0	0

Serpulidae	<i>Serpula</i> sp.	0.39	3.45	0	0.48	0.24	0	1.2	0	0
Spionidae	Spionidae indet.	6.73	8.62	64.29	1.44	10.14	0	0	0.4	0
Syllidae	<i>Brania pusilla</i>	1.78	0	0	0	2.66	6.93	2.41	0.4	1.67
	<i>Erinaceusyllis belizensis</i>	1.24	0	0	0	0.48	1.98	2.41	3.97	0
	<i>Eusyllis</i> sp.	0.23	0	0	0.96	0	0	0	0.4	0
	<i>Exogone breviantennata</i>	7.19	0	1.79	3.35	9.66	7.92	21.69	0	15.83
	<i>Exogone simplex</i>	3.48	0	0	0	0	2.97	14.46	0.79	23.33
	<i>Lamellisyllis</i> sp.	0.08	0	1.79	0	0	0	0	0	0
	<i>Opisthosyllis brunnea</i>	1.78	0	0	9.09	0.97	0	0	0	0
	<i>Parexogone anseforbansensis</i>	0.15	0	0	0	0.48	0	0	0	0
	<i>Perkinsyllis</i> sp.	0.08	0	0	0	0.24	0	0	0	0
	Syllidae indet. epit.	0.31	0	1.79	0	0	1.98	0	0.4	0
	Syllidae indet. juv.	5.1	1.72	0	7.18	0.97	0	3.61	15.08	4.17
	<i>Syllis</i> cf. <i>gerlachi</i>	2.24	0	0	0	0	3.96	0	8.73	2.5
	<i>Syllis</i> cf. <i>prolifera</i>	0.08	1.72	0	0	0	0	0	0	0
	<i>Syllis hyllebergi</i>	2.86	0	5.36	0	2.42	2.97	7.23	5.16	1.67
	<i>Syllis magellanica</i>	2.01	0	0	0	0.97	11.88	4.82	1.19	2.5
	<i>Syllis</i> sp.	9.59	1.72	5.36	4.31	4.35	0	1.2	35.71	1.67
	<i>Trypanosyllis auranticus</i>	1.78	10.34	1.79	1.91	1.21	0.99	0	2.38	0
	<i>Trypanosyllis</i> sp.1	1.08	0	0	0.96	0.24	2.97	0	2.38	1.67
	<i>Trypanosyllis</i> sp.2	2.09	0	0	0	0	0	0	10.71	0
	<i>Trypanosyllis zebra</i>	0.15	0	0	0	0.48	0	0	0	0
Terebellidae	Terebellidae indet.	4.49	0	1.79	6.7	10.39	0	0	0	0

Table S4. SIMPER results for the between-site contrast (Itaipu vs Sacristia). based on Bray–Curtis dissimilarities of $\log(x+1)$ abundances. The table lists taxa in decreasing order of contribution to the between-group dissimilarity and reports: Average (Bray) (mean contribution to D). % (percentage contribution). Cum % (cumulative percentage). and Mean (raw) — Itaipu / Sacristia (mean untransformed abundances per beach. averaged across replicates). Abbreviations: sp. = species undetermined; indet. = higher-rank identification; cf. = confer; juv. = juvenile; epit. = epitoke.

Species	Average (Bray)	%	Cum %	Mean (raw) Itaipu	Mean (raw) Sacristia
Orbiniidae	0.05311	6.7	6.7	4.5	1.67
Nereididae	0.05243	6.62	13.3	9.92	2.33
Spionidae	0.05003	6.31	19.6	7.17	0.08
<i>Exogone simplex</i>	0.04771	6.02	25.6	0	3.75
Sabellidae	0.04438	5.6	31.2	15.5	3.92
Cirratulidae	0.04317	5.45	36.7	1	2.5
<i>Syllis</i> sp	0.04314	5.44	42.1	2.58	7.75
<i>Exogone breviantennata</i>	0.04236	5.34	47.5	4	3.75
Syllidae juv.	0.03533	4.46	51.9	1.67	3.83
<i>Syllis hyllebergi</i>	0.0322	4.06	56	1.08	2
Terebellidae	0.03124	3.94	59.9	4.83	0
<i>Trypanosyllis auranticus</i>	0.02832	3.57	63.5	1.33	0.58
<i>Syllis magellanica</i>	0.02746	3.46	67	0.33	1.83
<i>Syllis</i> cf. <i>gerlachi</i>	0.02533	3.2	70.2	0	2.42

<i>Opisthosyllis brunnea</i>	0.02162	2.73	72.9	1.92	0
<i>Brania pusilla</i>	0.02109	2.66	75.6	0.92	1
<i>Pharyngocirrus gabriellae</i>	0.01701	2.15	77.7	0.25	0.25
<i>Trypanosyllis</i> sp1	0.01637	2.07	79.8	0.25	0.92
Eunicidae juv.	0.01523	1.92	81.7	0.25	0.33
<i>Erinaceusyllis belizensis</i>	0.01325	1.67	83.4	0.17	1.17
<i>Pseudoleiocyathella</i> cf. <i>fauveli</i>	0.01312	1.66	85	1.08	0
<i>Trypanosyllis</i> sp2	0.01277	1.61	86.6	0	2.25
<i>Serpula</i> sp	0.01025	1.29	87.9	0.33	0.08
<i>Lysidice unicornis</i>	0.01015	1.28	89.2	0	0.5
Arenicolidae indet.	0.00808	1.02	90.2	0	1.92
Syllidae epit.	0.00699	0.88	91.1	0.08	0.25
<i>Barantola</i> sp	0.00498	0.63	91.7	0.33	0
<i>Branchiomaldane</i> sp	0.00456	0.57	92.3	0	0.42
<i>Eunice</i> sp	0.00417	0.53	92.8	0.08	0.17
<i>Phragmatopoma</i> sp	0.00412	0.52	93.4	0.08	0.08
<i>Eusyllis</i> sp	0.00406	0.51	93.9	0.17	0.08
<i>Capitellethus</i> sp	0.00388	0.49	94.4	0.17	0
<i>Capitella</i> sp2	0.00353	0.45	94.8	0	0.25
<i>Leiocyathellides</i> cf. <i>analis</i>	0.00325	0.41	95.2	0.08	0
<i>Trypanosyllis zebra</i>	0.00314	0.4	95.6	0.17	0
<i>Capitella</i> sp1	0.00314	0.4	96	0.08	0.08
<i>Lamellisyllis</i> sp	0.00292	0.37	96.4	0.08	0
<i>Notomastus</i> sp	0.00285	0.36	96.7	0.08	0
<i>Scyphoproctus</i> sp	0.00285	0.36	97.1	0.08	0
<i>Lumbrineris</i> sp	0.00282	0.36	97.4	0.08	0
Capitellidae juvenil	0.00272	0.34	97.8	0.08	0
<i>Saccocirrus pussicus</i>	0.00272	0.34	98.1	0.08	0
<i>Syllis</i> cf. <i>prolifera</i>	0.00272	0.34	98.5	0.08	0
<i>Parheteromastides</i> sp	0.00261	0.33	98.8	0.08	0.08
<i>Lysidice</i> sp	0.00245	0.31	99.1	0	0.08
<i>Perkinsyllis</i> sp	0.00242	0.31	99.4	0.08	0
<i>Parexogone anseforbansensis</i>	0.00184	0.23	99.7	0.17	0
<i>Ophryotrocha</i> sp	0.00158	0.2	99.9	0.08	0
<i>Lysidice ninetta</i>	0.00116	0.15	100	0.08	0

Table S5. SIMPER results for the station contrast within Itaipu (Station 1 vs Station 2). based on Bray–Curtis dissimilarities of log(x+1) abundances. The table lists taxa in decreasing order of contribution to the between-group dissimilarity and reports: Average (Bray). %. Cum %. and Mean (raw) Station 1 / Station 2 (mean untransformed abundances per station. averaged across replicates). Abbreviations: sp. = species undetermined; indet. = higher-rank identification; cf. = confer; juv. = juvenile; epit. = epitoke.

Species	Average (Bray)	%	Cum %	Mean (raw) Station 1	Mean (raw) Station 2
Nereididae	0.08912	11.64	11.6	0.33	19.5
Terebellidae	0.06594	8.61	20.2	0.17	9.5

<i>Opisthosyllis brunnea</i>	0.05327	6.95	27.2	0	3.83
<i>Exogone breviantennata</i>	0.05153	6.73	33.9	0.17	7.83
Spionidae	0.05115	6.68	40.6	6.83	7.5
Syllidae juv.	0.04525	5.91	46.5	0.17	3.17
Sabellidae	0.04382	5.72	52.2	0.83	30.17
<i>Syllis</i> sp	0.03825	4.99	57.2	0.67	4.5
Orbiniidae	0.03668	4.79	62	5.33	3.67
<i>Pseudoleiocardia</i> cf. <i>fauveli</i>	0.03147	4.11	66.1	0	2.17
<i>Trypanosyllis auranticus</i>	0.02926	3.82	70	1.17	1.5
<i>Syllis hyllebergi</i>	0.02791	3.64	73.6	0.5	1.67
Cirratulidae	0.01847	2.41	76	0.33	1.67
<i>Brania pusilla</i>	0.01836	2.4	78.4	0	1.83
<i>Serpula</i> sp	0.01258	1.64	80	0.33	0.33
<i>Barantola</i> sp	0.01175	1.53	81.6	0	0.67
<i>Trypanosyllis</i> sp1	0.01111	1.45	83	0	0.5
<i>Syllis magellanica</i>	0.01111	1.45	84.5	0	0.67
Eunicidae juv.	0.0108	1.41	85.9	0.33	0.17
<i>Pharyngocirrus gabriellae</i>	0.00916	1.2	87.1	0.5	0
<i>Notomastus</i> sp	0.00742	0.97	88.1	0	0.17
<i>Scyphoproctus</i> sp	0.00742	0.97	89	0	0.17
<i>Trypanosyllis zebra</i>	0.0074	0.97	90	0	0.33
<i>Perkinsyllis</i> sp	0.0061	0.8	90.8	0	0.17
<i>Capitellethus</i> sp	0.00587	0.77	91.6	0.17	0.17
<i>Eusyllis</i> sp	0.00586	0.76	92.3	0	0.33
<i>Phragmatopoma</i> sp	0.00478	0.62	92.9	0	0.17
<i>Leiocardia</i> cf. <i>analis</i>	0.00458	0.6	93.5	0.17	0
<i>Lamellisyllis</i> sp	0.00427	0.56	94.1	0.17	0
Syllidae epit.	0.00427	0.56	94.7	0.17	0
<i>Lumbrineris</i> sp	0.00416	0.54	95.2	0.17	0
<i>Erinaceusyllis belizensis</i>	0.00415	0.54	95.7	0	0.33
<i>Parexogone anseforbansensis</i>	0.00415	0.54	96.3	0	0.33
Capitellidae juv.	0.00405	0.53	96.8	0.17	0
<i>Saccocirrus pussicus</i>	0.00405	0.53	97.3	0.17	0
<i>Syllis aff. prolifera</i>	0.00405	0.53	97.9	0.17	0
<i>Capitella</i> sp1	0.0037	0.48	98.4	0	0.17
<i>Ophryotrocha</i> sp	0.0037	0.48	98.8	0	0.17
<i>Eunice</i> sp	0.0037	0.48	99.3	0	0.17
<i>Parheteromastides</i> sp	0.00262	0.34	99.7	0	0.17
<i>Lysidice ninetta</i>	0.00262	0.34	100	0	0.17

Table S6. SIMPER results for the zone contrast within Sacristia (midlittoral vs upper infralittoral). based on Bray–Curtis dissimilarities of log(x+1) abundances. The table lists taxa in decreasing order of contribution to the between-group dissimilarity and reports: Average (Bray). %. Cum %. and Mean (raw) — midlittoral / upper infralittoral (mean untransformed abundances per zone. averaged across replicates).

Abbreviations: sp. = species undetermined; indet. = higher-rank identification; cf. = confer; juv. = juvenile; epit. = epitoke.

Species	Average (Bray)	%	Cum %	Mean (raw) midlittoral	Mean (raw) infralittoral
Cirratulidae	0.05913	7.85	7.8	1.5	3.5
<i>Exogone simplex</i>	0.05736	7.61	15.5	0.83	6.67
<i>Exogone breviantennata</i>	0.05501	7.3	22.8	1.33	6.17
<i>Syllis</i> sp	0.05338	7.08	29.8	15	0.5
Sabellidae	0.05209	6.91	36.8	5.33	2.5
Nereididae	0.04477	5.94	42.7	3.67	1
<i>Syllis</i> cf. <i>gerlachi</i>	0.04034	5.35	48	4.33	0.5
Syllidae juv.	0.03978	5.28	53.3	6.33	1.33
<i>Syllis hyllebergi</i>	0.03491	4.63	58	2.67	1.33
Orbiniidae	0.0349	4.63	62.6	1.5	1.83
<i>Syllis magellanica</i>	0.03483	4.62	67.2	2.5	1.17
<i>Trypanosyllis</i> sp2	0.02633	3.49	70.7	4.5	0
<i>Brania pusilla</i>	0.02566	3.41	74.1	1.33	0.67
<i>Trypanosyllis</i> sp1	0.02433	3.23	77.3	1.5	0.33
<i>Trypanosyllis auranticus</i>	0.02348	3.12	80.5	1.17	0
<i>Erinaceusyllis belizensis</i>	0.02152	2.86	83.3	2	0.33
Eunicidae juvenil	0.01698	2.25	85.6	0.17	0.5
<i>Pharyngocirrus gabriellae</i>	0.01641	2.18	87.7	0.33	0.17
Arenicolidae indet.	0.01547	2.05	89.8	0	3.83
<i>Lysidice unicornis</i>	0.0146	1.94	91.7	0.5	0.5
Syllidae epit.	0.00971	1.29	93	0.5	0
<i>Branchiomaldane</i> sp	0.00872	1.16	94.2	0	0.83
<i>Capitella</i> sp2	0.00727	0.97	95.1	0.5	0
<i>Serpula</i> sp	0.00654	0.87	96	0	0.17
<i>Eunice</i> sp	0.00576	0.77	96.8	0.33	0
<i>Lysidice</i> sp	0.00504	0.67	97.4	0.17	0
<i>Phragmatopoma</i> sp	0.00504	0.67	98.1	0.17	0
<i>Capitella</i> sp1	0.00364	0.48	98.6	0.17	0
Spionidae	0.00364	0.48	99.1	0.17	0
<i>Eusyllis</i> sp	0.00364	0.48	99.6	0.17	0
<i>Parheteromastides</i> sp	0.0033	0.44	100	0.17	0

Tabela S7. Environmental variables recorded at the rocky shore stations of Itaipu and Sacristia

Beach	Station	Temperature (C°)	Dissolved oxygen (mg/L)	Salinity (PSU)	pH
Itaipu	1	24.4	6.8	34.5	8.63
Itaipu	2	25.4	7.8	33.8	8.65
Sacristia	1	25.1	7.6	34.6	8.20
Sacristia	2	24.9	8.1	34.9	8.21