

Fisheries Science

Supplemental figure

Title

Interannual variability of larval fish assemblages associated with water masses in winter in the Taiwan Strait during 2007–2013

Authors

Yu-Kai Chen^{1,2}·Chia-Yi Pan²·Yi-Chen Wang¹·Hsiu-Ju Tseng²·Bo-Kun Su²·Ming-An Lee^{1,3*}

Author Affiliations

¹ Department of Environmental Biology and Fisheries Science, National Taiwan Ocean University, Keelung, Taiwan

² Fisheries Research Institute, Council of Agriculture, Keelung, Taiwan


³ Center of Excellence for Ocean Engineering, National Taiwan Ocean University, Keelung, Taiwan

*Corresponding author: Ming-An Lee

Address: No.2, Beining Rd., Jhongjheng District, Keelung City 202, Taiwan

Tel.: +886 24622192 ext. 5032

Email: malee@ntou.edu.tw



Online Resource Fig. S1 The Temperature–Salinity diagram (T-S diagram) for each cruise in the TS during the winters of 2007–2013. The area in the dotted lines denotes the water masses defined by Gong et al. (1996) and Ichikawa and Chaen (2000). KW, Kuroshio Water. The area between the solid lines denotes the water masses defined by Jan et al. (2006). KBW, Kuroshio Branch Water; NTMW, Northern Taiwan Mixed Water; MCCW, Mixed China Coastal Water. The stations with low temperature ($<16.5^{\circ}\text{C}$) and relatively low salinity (30.7–33.7psu) were regarded as CCC.