

Online Resource 3 Comparison of the age-standardised^a incidence rates of annual keratinocyte carcinomas in the Tasmanian Cancer Registry for Tasmania^b and in the National Non-Melanoma Skin Cancer Survey for Southern latitudes above 37°S [20]

	1985	1990	1995	2002
Tasmanian Cancer Registry				
Basal cell carcinoma				
Males	179 (163, 196)	291 (271, 311)	437 (414, 460)	560 (536, 585)
Females	85 (74, 97)	168 (153, 183)	256 (238, 274)	350 (331, 370)
Persons	129 (119, 139)	226 (213, 238)	339 (324, 353)	448 (432, 463)
Squamous cell carcinoma				
Males	78 (67, 89)	126 (114, 139)	231 (215, 248)	270 (253, 286)
Females	29 (23, 35)	46 (39, 53)	102 (92, 113)	143 (131, 155)
Persons	52 (46, 58)	82 (75, 89)	161 (152, 171)	202 (192, 212)
National Non-Melanoma Skin Cancer Survey				
Basal cell carcinoma				
Males	359 (223, 578)	421 (309, 532)	511 (398, 659)	646 (505, 827)
Females	307 (182, 519)	230 (151, 309)	360 (267, 486)	462 (349, 614)
Persons	333 (234, 474)	323 (256, 390)	430 (355, 522)	547 (454, 659)
Squamous cell carcinoma				
Males	64 (21, 199)	80 (35, 125)	229 (157, 334)	306 (217, 433)
Females	43 (11, 174)	13 (0, 30)	114 (66, 197)	171 (109, 267)
Persons	53 (22, 127)	47 (23, 71)	167 (123, 228)	232 (177, 306)

^aStandardised to the 1960 World population.

^bTasmania is located between 40°S and 43°S.