

Excluding external iliac node irradiation during neoadjuvant radiotherapy decreases lower intestinal toxicity without compromising efficacy in T4b rectal cancer patients with tumours involving the anterior structures

Authors: Anchuan Li^{1,2,3,4}, Miaobin Mao^{1,3,4}, Runfan Chen¹, Pan Chi⁵; Ying Huang⁵; Benhua Xu^{1,2,3,4*}

¹ Department of Radiation Oncology, Fujian Medical University Union Hospital, Fuzhou, 350001, China;

² Department of Radiation Oncology, College of Clinical Medicine, Fujian Medical University, Fuzhou, 350001, China;

³ Fujian Key Laboratory of Intelligent Imaging and Precision Radiotherapy for Tumors, Fujian Medical University, Fuzhou, 350001, China;

⁴ Clinical Research Center for Radiology and Radiotherapy of Fujian Province (Digestive, Hematological and Breast Malignancies), Fuzhou, 350001, China;

⁵ Department of Gastrointestinal Surgery, Fujian Medical University Union Hospital,

Fuzhou, 350001, China;

* **Corresponding author:** Ben-hua Xu, Department of Radiation Oncology, Fujian Medical University Union Hospital, Xinquan Road 29, Fuzhou, 350001, China; Tel:

(86) 0591-86218731; Fax: (86) 0591-86218731; Email: benhuaxu@163.com.




Figure S1 Kaplan - Meier survival curves of patients with cN positivity treated with and without EIN irradiation. A) locoregional recurrence-free survival; B) distant metastasis-free survival; C) overall survival; D) progression-free survival




Figure S2 Schematic diagram of sites of locoregional recurrence