

## **Supplementary Material**

# **Neutrophil Extracellular Traps (NETs) are associated with Type 2 Diabetes and Diabetic Foot Ulcer Related Amputation: A Prospective Cohort Study**

**Irshat Ibrahim<sup>1†</sup>, Yilimire Nuermaiti<sup>2†</sup>, Gulijianaiti Maimaituxun<sup>3†</sup>, Xinling Luo<sup>1</sup>, Mailudemu Maimaituxun<sup>2</sup>, Azimat Akbar<sup>1</sup>, Kahaer Tuerxun<sup>1\*</sup>, Yuanquan Wu<sup>1\*</sup>**

<sup>1</sup> Department of General Surgery, The First People's Hospital of Kashi Prefecture, Kashi 844000, China.

<sup>2</sup> Department of Endocrinology, The First People's Hospital of Kashi Prefecture, Kashi 844000, China.

<sup>3</sup> Department of Neurology, The First People's Hospital of Kashi Prefecture, Kashi 844000, China.

† These authors contributed equally to this work and share first authorship.

### **\* Correspondence:**

Yuanquan Wu: doctor\_wuyq@163.com; Kahaer Tuerxun: abdoctor@163.com

**Supplementary Figure S1.** Plots demonstrating the Spearman correlation for NETs (MPO-DNA complex) levels and HbA1c in (A) DFU group and (B) healthy controls, with Spearman correlation coefficient of 0.123 and -0.128 respectively without statistically significant differences.

