

SUPPLEMENTARY MATERIAL

Ruxolitinib Cream As-Needed Monotherapy Demonstrates Sustained Disease and Symptom Control in Patients with Mild to Moderate Atopic Dermatitis: Pooled Analysis from Two Phase 3 Studies


Andrew Blauvelt, MD, MBA,^{1,2} Leon Kircik, MD,^{3,4} Eric L. Simpson, MD, MCR,⁵ Peter Lio, MD,⁶ Lawrence F. Eichenfield, MD,⁷ Vimal H. Prajapati, MD,⁸⁻¹³ Parbeer Grewal, MD,¹⁴ Charles Lynde, MD,^{15,16} H. Chih-ho Hong, MD,^{17,18} Daniel Sturm, PharmD,¹⁹ Howard Kallender, PhD,¹⁹ Haobo Ren, PhD,¹⁹ Dilan Paranagama, PhD,¹⁹ Grace K. W. Wong, PhD,²⁰ Bob Geng, MD⁷

¹Oregon Medical Research Center, Portland, OR, USA; ²Blauvelt Consulting, LLC, Annapolis, MD, USA; ³Icahn School of Medicine at Mount Sinai, New York, NY, USA; ⁴Indiana University School of Medicine, Indianapolis, IN, USA; ⁵Oregon Health & Science University, Portland, OR, USA; ⁶Feinberg School of Medicine, Northwestern University, Chicago, IL, USA; ⁷University of California San Diego, La Jolla, CA, USA; ⁸Division of Dermatology, Department of Medicine, University of Calgary, Calgary, AB, Canada; ⁹Section of Community Pediatric, Department of Pediatrics, University of Calgary, Calgary, AB, Canada; ¹⁰Section of Pediatric Rheumatology, Department of Pediatrics, University of Calgary, Calgary, AB, Canada; ¹¹Dermatology Research Institute, Calgary, AB, Canada; ¹²Skin Health & Wellness Centre, Calgary, AB, Canada; ¹³Probit Medical Research, Calgary, AB, Canada; ¹⁴Rejuvenation Medical, Edmonton, AB, Canada; ¹⁵Lynde Institute for Dermatology, Markham, ON, Canada; ¹⁶Probit Medical Research, Markham, ON, Canada; ¹⁷Department of Dermatology and Skin Science, University of British Columbia, Vancouver, BC, Canada; ¹⁸Probit Medical Research, Surrey, BC, Canada; ¹⁹Incyte Corporation, Wilmington, DE, USA; ²⁰Incyte Biosciences Canada, Pointe-Claire, QC, Canada

Corresponding author: Bob Geng, MD

University of California San Diego
3030 Children's Way # 2
La Jolla, CA, USA. 92123.
Phone: 858-966-5961
Email: geng.bob@gmail.com


Supplementary Material Fig. S1. Percentage of Visits* With IGA 0/1 Among Patients Who Achieved or Did Not Achieve IGA-TS, EASI-75, or Itch NRS 0/1 at Week 8



* Included patients with ≥ 2 visits.

EASI-75, $\geq 75\%$ improvement from baseline in Eczema Area and Severity Index; IGA, Investigator's Global Assessment; IGA-TS, IGA treatment success; LTS, long-term safety; NRS, numerical rating scale.


Supplementary Material Fig. S2. Percentage of Visits* With IGA 0/1 in the LTS Period
According to the Number of Prior Lines of Therapy



* For patients with ≥ 2 visits.

IGA, Investigator's Global Assessment; LTS, long-term safety.


Supplementary Material Fig. S3. Percentage of Treatment-Free Days Between Study Visits for Patients Who Did or Did Not Achieve (A) IGA-TS, (B) EASI-75, or (C) Itch NRS 0/1 at Week 8



* Line corresponds to affected BSA in the last week of the stated range.


BSA, body surface area; EASI-75, $\geq 75\%$ improvement from baseline in Eczema Area and Severity Index; IGA, Investigator's Global Assessment; IGA-TS, IGA treatment success; LTS, long-term safety; NRS, numerical rating scale.

Supplementary Material Fig. S4. Time Off Treatment by Baseline Disease Severity and Extent



* As median percentage of the entire long-term safety period; 244 total patients were included in this analysis. BSA, body surface area; IGA, Investigator's Global Assessment.

Supplementary Material Fig. S5. Time Off Treatment by Number of Prior Lines of Therapy



* As median percentage of the entire long-term safety period.