

Real-World Ritlecitinib Treatment of Severe Alopecia Areata in the United States: Patient Characteristics and Physician Satisfaction According to a Secondary Database Analysis


Samantha K. Kurosky¹, Ashley S. Cha-Silva¹, Jenny Austin², Arash Mostaghimi³, Chesahna Kindred^{4,5}, Genevieve Gauthier⁶, Mojgan Sadrarhami¹, Peter Anderson², Benjamin Ungar⁷

1. Pfizer Inc, New York, NY, USA
2. Adelphi Real World, Bollington, UK
3. Brigham and Women's Hospital, Boston, MA, USA
4. Kindred Hair & Skin Center, Columbia, MD, USA
5. Howard University, Washington, DC, USA
6. Pfizer Inc, Kirkland, QC, Canada
7. Mount Sinai, New York, NY, USA

Corresponding author: Samantha K. Kurosky (Samantha.Kurosky@pfizer.com)


SUPPLEMENTARY MATERIAL

Figure S1. Study design



AA, alopecia areata; DSP, disease specific programme.


Figure S2. Current disease severity* by ritlecitinib treatment duration among patients[†] who had (A) severe or (B) very severe disease immediately prior to ritlecitinib initiation



*Disease severity currently and immediately prior to ritlecitinib initiation were based on physician assessment.

[†]Patients with a known duration of ritlecitinib treatment.

Figure S3. Treatment goals* among patients receiving ritlecitinib (N=123) (A) and reasons for selecting ritlecitinib* among patients receiving ritlecitinib (n=113) (B)



*Reported for $\geq 10\%$ of patients.