

Extended Depth of Field Intraocular Lens for Cataract After Vitrectomy for Retinal Detachment: A Prospective Trial

Efstathios Vounotrypidis¹, Dorian Begaj¹, Rupert Kamnig¹, Julie Meyer¹, Melih Parlak¹, Christian M. Wertheimer¹, Armin Wolf¹

1. Department of Ophthalmology, University Hospital Ulm, Ulm, Germany

Corresponding author:

Prof. Dr. med. Armin Wolf, FEBO

Department of Ophthalmology,

University Hospital Ulm,

Prittwitzstr. 43,

D-89075, Ulm, Germany


Phone: +4973150059004

Fax: +4973150059002

Email: armin.wolf@uniklinik-ulm.de


Prior presentation: Parts of this work were presented at the 42nd ESCRS congress in Barcelona, Spain and the FEOph Symposium DOG 2024 in Berlin, Germany.

Supplemental Material 1:




Percentage of patients with regard to their satisfaction with their vision.

Supplemental Material 2:



Results of the German Catquest-9SF questionnaire for specific daily life activities in patients after unilateral and bilateral surgery.

Supplemental Material 3:



Spectacle independence through intraocular lens satisfaction questionnaire (IOLSAT) questionnaire in bright (BL) and dim light (DL) conditions at 6 months after implantation of IOL after unilateral and bilateral surgery.

“In the past 7 days, how often did you need to wear eyeglasses to see in bright light/dim light conditions”? (0 = never; 1 = rarely; 2 = sometimes; 3 = most of the time; 4 = all the time).

Percentages of patients who answered “never” or “rarely” are highlighted.