

Article title: Prevalence and risk factors for psychotropic medication use in older adults in Australia: a nationwide, data linkage study

Journal: Drugs & Aging

Hieu T. Le^a, Edward C.Y. Lau^a, Weisi Chen^a, Christine Y. Lu^{a, b, c}, Tuan A. Nguyen^{d, e}, Lee-Fay Low^f, Sarah N Hilmer^b, Yun-Hee Jeon^g and Edwin C.K. Tan^{a, b*}

^a*The University of Sydney School of Pharmacy, Faculty of Medicine and Health, The University of Sydney, Sydney, New South Wales, Australia*

^b*Kolling Institute, Faculty of Medicine and Health, The University of Sydney and the Northern Local Health District, Sydney, New South Wales, Australia*

^c*Department of Pharmacy, Royal North Shore Hospital, St Leonards, New South Wales, Australia*

^d*Division of Social Gerontology, National Ageing Research Institute, Parkville, Victoria, Australia.*

^e*School of Health Sciences, Swinburne University of Technology, Hawthorn, Victoria, Australia*

^f*The University of Sydney School of Health Sciences, Faculty of Medicine and Health, The University of Sydney, Sydney, New South Wales, Australia*


^g*Susan Wakil School of Nursing and Midwifery, Faculty of Medicine and Health, The University of Sydney, Sydney, New South Wales, Australia*

***Corresponding author:**








Dr Edwin C.K. Tan

Post: The University of Sydney School of Pharmacy, Faculty of Medicine and Health, Pharmacy and Bank Building A15, The University of Sydney, NSW, Australia, 2006

Email: edwin.tan@sydney.edu.au



Supplementary Figure 1: Study flow chart



Supplementary Figure 2: Prevalence of psychotropic medication use by age group and sex