

AGGRESSIVE SUBTYPES OF LARYNGEAL CHONDROSARCOMA AND THEIR CLINICAL BEHAVIOR – A SYSTEMATIC REVIEW.

Cesare Piazza ^{1,2}, Claudia Montenegro ^{1,2}, Michele Tomasoni ¹, Ilmo Leivo ^{3,4}, Göran Stenman ⁵, Abbas Agaimy ⁶, Roderick H.W. Simpson ⁷, Nina Zidar ⁸, Alfio Ferlito ⁹.

¹Unit of Otorhinolaryngology – Head and Neck Surgery, ASST Spedali Civili of Brescia, Brescia, Italy

²Department of Medical and Surgical Specialties, Radiological Sciences, and Public Health, University of Brescia, School of Medicine, Brescia, Italy

³Institute of Biomedicine, Pathology, University of Turku

⁴Turku University Central Hospital, Turku

⁵Sahlgrenska Center for Cancer Research, Department of Pathology, University of Gothenburg, Gothenburg, Sweden

⁶Institute of Pathology, University Hospital Erlangen, Friedrich-Alexander University Erlangen-Nürnberg, Erlangen, Germany

⁷Department of Anatomical Pathology, University of Calgary, Calgary, AB, Canada

⁸Institute of Pathology, Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia

⁹Coordinator of the International Head and Neck Scientific Group, Padua, Italy

This article was written by members and invitees of the International Head and Neck Scientific Group (www.IHNSG.com)

The Authors have no conflict of interest related to this manuscript

Corresponding author:

Claudia Montenegro, MD

Unit of Otorhinolaryngology – Head and Neck Surgery

ASST Spedali Civili of Brescia/ University of Brescia, School of Medicine

Brescia, Italy

Email address: c.montenegro@unibs.it

SUPPLEMENTARY TABLE

Table S1 – Quality assessment of case series included in the present systematic review (N=9).

[illegible]