

Supplementary Table 1. Laboratory Data

Variable	Admission	Discharge	Reference Range, Adults
Blood			
Hematocrit (%)	35	36.1	36–46
Hemoglobin (g/dL)	11.7	11.8	12–16
Red-cell count (per μ L)	3,740,000	3,790,000	4,200,000–5,400,000
Mean corpuscular volume (fl)	94.7	95.3	81–94
Red blood cell distribution width (%)	12.8	13.3	11.8–14.8
White-cell count (per μ L)	13100	9060	4000–10000
Differential count			
Neutrophils (per μ L)	10,320	6480	1500–7500
Lymphocytes (per μ L)	1,950	1860	1000–4800
Monocytes (per μ L)	790	700	300–1200
Eosinophils (per μ L)	10	10	0–700
Basophils (per μ L)	30	20	0–200
Platelet count (per μ L)	308,000	289,000	140,000–440,000
Prothrombin time (sec)	11.7	12.3	9.5–12
Prothrombin time (%)	95	89	70–130
Prothrombin-time international normalized ratio	1.0	1.09	0.8–1.2
Activated-partial thromboplastin time (sec)	22.5	26.9	22–38
Activated-partial thromboplastin time (ratio)	0.76	0.91	0.73–1.27
Sodium (mEq/L)	140	139	135–145
Potassium (mEq/L)	4.5	3.9	3.5–5.1
Chloride (mEq/L)	103	100	95–110
Calcium (mg/dL)	8.9	8.6	8.6–10.2
Magnesium (mg/dL)	1.8	1.7	1.7–2.5
Creatinine (mg/dL)	0.66	0.58	0.5–1.10
Glucose (mg/dL)	100	97	65–110
Fibrinogen (mg/dL)	196	216	200–400
Alanine aminotransferase (U/L)	28	48	10–35
Total bilirubin	0.4	0.4	0.2–1
Lactate dehydrogenase (U/L)	163	203	135–214
C-reactive protein (mg/L)	<5	<5	<5
Total IgE (KU/L)	359	82.8	0–85

Variable	Admission	Reference Range, Adults
Eosinophil cationic protein (μ g/L)	11	0–15
Basal serum tryptase (μ g/L)	3.5	0–13.4
NT-proBNP (pg/mL)	65	1–125
Erythrocyte sedimentation rate (mm/hr)	3	2–30
Procalcitonin (ng/mL)	0.06	<0.5
Troponin T (pg/mL)	<3	<14
Protein		
Total (g/dL)	6.7	6–8.2
Albumin (%)	58.4	55.8–66.1
α 1-globulins (%)	3.4	2.9–4.9
α 2-globulins (%)	11.6	7.1–11.8
β 1-globulins (%)	5.4	4.7–7.2

β2-globulins (%)	4.8	3.2–6.5
γ-globulins (%)	16.4	11.1–18.8
IgG (mg/dL)	1180	700–1600
IgA (mg/dL)	271	70–400
IgM (mg/dL)	69	40–230
IgG1 (mg/dL)	750	490–1140
IgG2 (mg/dL)	357	150–640
IgG3 (mg/dL)	79	20–110
IgG4 (mg/dL)	35	8–82
Complement C4 (g/L)	0.13	0.1–0.4
Antinuclear antibodies (IFI on HEp2 cells)	1:80; speckled pattern	1:80
ENA CTD-screen (ratio)	0.2	0–1
Antibodies to anti-neutrophil cytoplasm (ANCA)	Not present	Not present
anti-PR3 antibodies (UI/mL)	0.3	0–3
anti-MPO antibodies (UI/mL)	0.2	0–5
Anti-Respiratory Syncytial Virus antibodies (IgA)	Not present	Not present
Anti-Parainfluenza virus 1-2-3 antibodies	Not present	Not present
Galactomannan antigen	Negative	Negative
1-3-β-D-Glucan (pg/mL)	Negative	Negative
Influenza Virus A (PCR)	Negative	Negative
Influenza Virus B (PCR)	Negative	Negative
SARS-CoV-2 (PCR)	Negative	Negative
Soluble <i>Legionella pneumophila</i> antigens	Not present	Not present
Soluble <i>Streptococcus pneumoniae</i> antigens	Not present	Not present
IgG Anti- <i>Strongyloides</i> antibodies	Not present	Not present
IgG anti- <i>Toxocara</i> antibodies	Not present	Not present
Ova and parasite stool test	Negative	–
Fecal cultures	Negative	–
Anisakis-specific IgE (KU/L)	6.98	<0.10
Velvet grass-specific IgE (KU/L)	5.25	<0.10
Wall pellitory-specific IgE (KU/L)	2.98	<0.10
Cat dander-specific IgE (KU/L)	0.52	<0.10
House dust mite-specific IgE (KU/L)	0.22	<0.10
Milk-specific IgE (KU/L)	0.17	<0.10
Egg white-specific IgE (KU/L)	0.11	<0.10
Shrimp-specific IgE (KU/L)	0.02	<0.10
Latex-specific IgE (KU/L)	0.02	<0.10
rBet v 1 (KU/L)	0.01	<0.10
rBet v 2 (KU/L)	0.02	<0.10
rPru p 3 (KU/L)	0.03	<0.10
rTri a 19 (KU/L)	0.02	<0.10
rTri a 14 (KU/L)	0.01	<0.10
Alpha-gal (KU/L)	0	<0.10
Arterial blood gases		
Fraction of inspired oxygen (%)	21	
pH	7.44	7.35–7.45
PaCO ₂ (mm Hg)	37	32–45
PaO ₂ (mm Hg)	91	83–108

Urine		
Color	Yellow	Yellow
Clarity	Clear	Clear
pH	6	5.0–6.5
Specific gravity	1.023	1.015–1.025
Protein	Negative	Negative
Glucose	Negative	Negative
Ketones	Negative	Negative
Bilirubin	Negative	Negative
Nitrite	Negative	Negative
Leukocyte esterase	250	Negative
Red cells (per μL)	10	<20
Leukocytes (per μL)	127	<18