

Assessing hoverfly (Diptera: Syrphidae) assemblages in the Cerrado biome: insights from different vegetation types, climate variables, and collection techniques

Anna Mara Ferreira Maciel^{1,2*} ([0000-0002-7735-9332](#)), Mírian Nunes Morales²([0000-0003-2114-8656](#)), Lucas Del Bianco Faria³([0000-0002-7230-800X](#)), Marcel Gustavo Hermes^{2,4}([0000-0002-9322-4518](#))

***Corresponding author:**

¹ Department of Entomology and Acarology, Escola Superior de Agricultura “Luiz de Queiroz”, Universidade de São Paulo, Piracicaba, 13418-900, SP, Brazil.

² Department of Entomology, Escola de Ciências Agrárias de Lavras, Universidade Federal de Lavras, Lavras, 37200-900, MG, Brazil.

annamaramaci@gmail.com.

Supplementary Information – Online Resource 3

Supplementary Figure 2 Interpolation and extrapolation curves of Syrphidae diversity based on Hill numbers for sampling methods. (a) Species richness ($q = 0$), (b) Shannon diversity expressed as true diversity ($q = 1$), and (c) Simpson diversity expressed as true diversity ($q = 2$). Solid lines represent interpolation and dashed lines represent extrapolation; shaded areas indicate 95% confidence intervals. Curves were generated using standardized sampling effort.

