








Control without LPS or THP-1 supernatant	Control with 5 µg/ml LPS	Control with 50 µg/ml LPS	Control with supernatant from THP-1 cells
			



Cells pre-incubated with 5 µg/ml LPS			
Compounds	10 µM	50 µM	100 µM
TP1			
TP4			
TP5			

Cells pre-incubated with 5 µg/ml LPS			
Compounds	10 µM	50 µM	100 µM
TP6			
TP7			
TP8			



Cells pre-incubated with 5 µg/ml LPS			
Compounds	10 µM	50 µM	100 µM
TP9			
TP10			


Cells pre-incubated with 50 µg/ml LPS			
Compounds	10 µM	50 µM	100 µM
TP1			
TP4			
TP5			

Cells pre-incubated with 50 µg/ml LPS			
Compounds	10 µM	50 µM	100 µM
TP6			
TP7			
TP8			

Cells pre-incubated with 50 µg/ml LPS			
Compounds	10 µM	50 µM	100 µM
TP9			
TP10			

Cells pre-incubated with THP-1 supernatant			
Compounds	10 μ M	50 μ M	100 μ M
TP1			
TP4			
TP5			

Cells pre-incubated with THP-1 supernatant			
Compounds	10 μ M	50 μ M	100 μ M
TP6			
TP7			
TP8			

Cells pre-incubated with THP-1 supernatant			
Compounds	10 μ M	50 μ M	100 μ M
TP9			
TP10			