

Supplementary materials.

Table S1. Secondary structure composition of tuber powder proteins.

Band (cm ⁻¹)	center (cm ⁻¹)	Band assignment	Secondary structure type	References
1606	1606		β -sheet	[1]
1628	[1620 -1644]		β -sheet	[2, 3]
1645	[1640-1648]		Random coil	[3]
1659	[1649-1659]		α-helix	[4]
1673	[1660-1688]		β-turn	[3, 5]

References

- [1] Agnieszka S, Magdalena R, Jan B, Katarzyna W, Malgorzata B, Krzysztof H, Danuta K. Phytotoxic Effect of Fiber Hemp Essential Oil on Germination of Some Weeds and Crops. Essential Oil-Bear Plants. 2016; 19: 262–276. <https://doi.org/10.1080/0972060X.2015.1137236>.
- [2] Bock JK, Damodaran S. Bran-induced changes in water structure and gluten conformation in model gluten dough studied by Fourier transform infrared spectroscopy. Food Hydrocoll. 2013; 31: 146–155. <https://doi.org/10.1016/j.foodhyd.2012.10.014>
- [3] Kong J, Yu S. Fourier Transform Infrared Spectroscopic Analysis of Protein Secondary Structures. Acta. Biochim. Biophys. Sin. 2007; 39: 549–559. <https://doi.org/10.1111/j.1745-7270.2007.00320.x>
- [4] Van Velzen EJJ, Van Duynhoven JPM, Pudney P, Weegels PL., Van der Maas JH, Cereal Chem. 80, 378–382 (2003).<https://doi.org/10.1094/CCHEM.2003.80.4.378>
- [5] Yang H, Yang S, Kong J, Dong A, Yu S. Obtaining information about protein secondary structures in aqueous solution using Fourier transform IR spectroscopy Nat. Protoc. 2015 ; 10, 382–396. <https://doi.org/10.1038/nprot.2015.024>