

Supplementary Table 6. Effects of Free Radical Protective Compounds (100 μ M) on SIRT1 Rate.

Compound	Ratio to Control Rate Mean $\pm$ SE	Replicates	Protective Mechanism
Hinokitiol (b-Thujaplicin; 2-hydroxy-4-isopropyl-2,4,6-cycloheptatrien-1-one)	2.48 $\pm$ 0.15	2	Iron Chelator
L-(+)-Ergothioneine ((S)-a-Carboxy-2,3-dihydro-N,N,N-trimethyl-2-thioxo-1H-imidazole-4-ethanaminium inner salt)	2.06 $\pm$ 0.48	2	Antioxidant, Peroxynitrite Scavenger
Caffeic Acid Phenyl Ester	1.80 $\pm$ 0.16	2	Iron Chelator
MCI-186 (3-Methyl-1-phenyl-2-pyrazolin-5-one)	1.2513 $\pm$ 0.0080	2	Radical Scavenger and Antioxidant
HBED (N,N'-Di-(2-hydroxybenzyl)ethylenediamine-N,N'-diacetic acid•HCl•H ₂ O)	1.150 $\pm$ 0.090	2	Iron Chelator
Ambroxol (trans-4-(2-Amino-3,5-dibromobenzylamino)cyclohexane•HCl)	1.075 $\pm$ 0.0026	2	Radical Scavenger
U-83836E ((-)-2-((4-(2,6-di-1-Pyrrolidinyl-4-pyrimidinyl)-1-piperazinyl)methyl)-3,4-dihydro-2,5,7,8-tetramethyl-2H-1-benzopyran-6-ol•2HCl)	1.030 $\pm$ 0.055	2	"Lazaroid" aminosteroid, Peroxidation inhibitor
Trolox (6-Hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid)	0.995 $\pm$ 0.019	2	Antioxidant

Abbreviation: SE, Standard error of the mean. Rate measurements with 25 μ M NAD⁺ and 25 μ M p53-382 acetylated peptide substrate were performed as described in Methods. All ratio data were calculated from experiments in which the total deacetylation in the control reaction was 0.25-1.25 μ M peptide or 1-5% of the initial concentration of acetylated peptide.