

Acknowledgements The authors thank S. C. Leung and M. Kokubo for helpful discussions. We also thank the staff at the Southern African Large Telescope, the Gemini-North telescope, the Nordic Optical Telescope, the Isaac Newton Telescope, the Liverpool Telescope and the Kiso Schmidt telescope for observations and people who carried out follow-up observations which were unfruitful owing to the weather. Simulations for the He detonation models were carried out on a Cray XC30 at the Center for Computational Astrophysics, National Astronomical Observatory of Japan. The work is partly supported by the World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan, Grants-in-Aid for Scientific Research of JSPS (16H01087 and 26287029 for M.D. and J.J.; 26800100 and 17H02864 for K.M.; 16H06341, 16K05287, and 15H02082 for T.S.; 26400222, 16H02168, and 17K05382 for K.N.; 15H02075, 16H02183, and 17H06363 for M.T.; 15H05892 for S.M.) and the research grant program of the Toyota foundation (D11-R-0830). S.W.J. acknowledges support from the US National Science Foundation through award AST-1615455. M.D.S acknowledges generous support provided by the Danish Agency for Science and Technology and Innovation realized through a Sapere Aude Level 2 grant, the Instrument-center for Danish Astrophysics (IDA), and by a research grant (13261) from VILLUM FONDEN. The Hyper Suprime-Cam (HSC) collaboration includes the astronomical communities of Japan and Taiwan, and Princeton University. The HSC instrumentation and software were developed by the National Astronomical Observatory of Japan (NAOJ), the Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU), the University of Tokyo, the High Energy Accelerator Research Organization (KEK), the Academia Sinica Institute for Astronomy and Astrophysics in Taiwan (ASIAA), and Princeton University.

Funding was contributed by the FIRST program from Japanese Cabinet Office, the Ministry of Education, Culture, Sports, Science and Technology (MEXT), the Japan Society for the Promotion of Science (JSPS), Japan Science and Technology Agency (JST), the Toray Science Foundation, NAOJ, Kavli IPMU, KEK, ASIAA, and Princeton University. The Pan-STARRS1 Surveys (PS1) have been made possible through contributions of the Institute for Astronomy, the University of Hawaii, the Pan-STARRS Project Office, the Max-Planck Society and its participating institutes, the Max Planck Institute for Astronomy, Heidelberg and the Max Planck Institute for Extraterrestrial Physics, Garching, The Johns Hopkins University, Durham University, the University of Edinburgh, Queen's University Belfast, the Harvard-Smithsonian Center for Astrophysics, the Las Cumbres Observatory Global Telescope Network Incorporated, the National Central University of Taiwan, the Space Telescope Science Institute, the National Aeronautics and Space Administration under Grant No. NNX08AR22G issued through the Planetary Science Division of the NASA Science Mission Directorate, the National Science Foundation under Grant No. AST-1238877, the University of Maryland, and Eotvos Lorand University (ELTE). This paper makes use of software developed for the Large Synoptic Survey Telescope. We thank the LSST Project for making their code available as free software at <http://dm.lsst.org>. This work is also based on observations obtained at the Gemini Observatory (program: GN-2016A-DD-7), which is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the National Research Council (Canada), CONICYT (Chile), Ministerio de Ciencia, Tecnología e Innovación Productiva (Argentina), and Ministério da Ciência, Tecnologia e Inovação (Brazil).