

Single-cell transcriptional changes associated with drug tolerance and response to combination therapies in cancer

Alexandre F. Aissa¹, Abul B.M.M.K. Islam^{1,2}, Majd M. Ariss¹, Cammille C. Go¹, Alexandra E. Rader¹, Ryan D. Conrardy¹, Alexa M. Gajda¹, Carlota Rubio-Perez³, Klara Valyi-Nagy⁴, Mary Pasquinelli⁵, Lawrence E. Feldman⁵, Stefan J. Green⁶, Nuria Lopez-Bigas³, Maxim V. Frolov¹ and Elizaveta V. Benevolenskaya^{1*}.

¹ Department of Biochemistry and Molecular Genetics, University of Illinois at Chicago 900 S. Ashland Ave, Chicago 60607, USA

² Department of Genetic Engineering and Biotechnology, University of Dhaka, Dhaka 1000, Bangladesh

³ Biomedical Genomics Lab, Institute for Research in Biomedicine (IRB), Barcelona 08003, Spain



⁴ Department of Pathology, University of Illinois at Chicago, 840 South Wood Street, Chicago, IL, 60612, USA

⁵ Department of Medicine, Section of Hematology/Oncology, University of Illinois at Chicago, 840 South Wood Street, Chicago, IL, 60612, USA


⁶ Genome Research Core, Research Resources Center, University of Illinois at Chicago

* Corresponding author, e-mail: evb@uic.edu


Source Data



Source Data 1. Immunoblot analysis of cell lysates prepared from PC9 cells using antibodies to TACSTD2.



Source Data 2. Vinculin used as a loading control for the TACSTD2 experiment.



* Anti-rabbit IgG for membrane ② was done in 10.31.19
* Better exposure for membrane ① is on the back.

Source Data 3. Immunoblot analysis of cell lysates prepared from PC9 cells using antibodies to SERPINE1.

