

Changes in event soil moisture-temperature coupling may intensify very extreme heat beyond expectations

Supplementary information

Douglas Maraun¹, Reinhard Schiemann², Albert Osso¹ and Martin Jury¹

¹ Wegener Center for Climate and Global Change,
University of Graz, Graz, Austria

² National Centre for Atmospheric Research,
Department of Meteorology, University of Reading, Reading, UK




Figure S 1: Daily maximum temperature and its changes for ACCESS-ESM1-5. **a** Present day (1990-2014) warm season (annual mean for 20S to 20N) daily maximum temperature and **b** its projected changes (2076-2100 vs. 1990-2014) under the SSP5-8.5 scenario. Corresponding present day values and changes for the 2-year (**c,d**) and 200-year (**e,f**) return levels of daily maximum temperature T_X .




Figure S 2: Daily maximum temperature and its changes for MIROC6. a Present day (1990-2014) warm season (annual mean for 20S to 20N) daily maximum temperature and b its projected changes (2076-2100 vs. 1990-2014) under the SSP5-8.5 scenario. Corresponding present day values and changes for the 2-year (c,d) and 200-year (e,f) return levels of daily maximum temperature T_X .




Figure S 3: Difference in changes of moderate to extreme temperature return levels. Changes (2076-2100 vs. 1990-2014) in 5- (a-c), 10- (d-f), 20- (g-i), 50- (j-l) and 100-year (m-o) return levels minus changes in 2-year return levels of daily maximum temperature T_X for MPI-ESM1-2-LR (a,d,g,j,m), ACCESS-ESM1-5 (b,e,h,k,n) and MIROC6 (c,f,i,l,o). Differences in hatched regions are not significant at the 95% level (see Methods for detail).




Figure S 4: Present day seasonal mean coupling. a MPI-ESM1-2-LR, b ACCESS-ESM1-5 and c MIROC6.




a**b****c**

Figure S 5: Differences in changes between moderate and mean heat events
 Difference in changes (2076-2100 vs. 1990-2014) of 2-year return values and seasonal mean of daily maximum temperature T_X for **a** MPI-ESM1-2-LR, **b** ACCESS-ESM1-5 and **c** MIROC6 under the SSP5-8.5 scenario.




Figure S 6: Driving processes for ACCESS-ESM1-5. **a** Present (1990-2014) soil moisture temperature coupling strength (strong negative values indicate strong coupling), **b** present (1990-2014) event soil moisture, **c** event soil moisture changes **d** and warm season (annual mean for 20S to 20N) precipitation changes (2076-2100 vs. 1990-2014).




Figure S 7: Driving processes for MIROC6. **a** Present (1990-2014) soil moisture temperature coupling strength (strong negative values indicate strong coupling), **b** present (1990-2014) event soil moisture (note the different color scale compared to the two other models), **c** event soil moisture changes **d** and warm season (annual mean for 20S to 20N) precipitation changes (2076-2100 vs. 1990-2014).




Figure S 8: Difference in changes of extreme temperature return levels for near future projections for ACCESS-ESM1-5. Difference between changes in 200-year and 2-year return levels of daily maximum temperature T_X for near future climate (2036-2060 vs. 1990-2014).




Figure S 9: Difference in changes of extreme temperature return levels for near future projections for MIROC6. Difference between changes in 200-year and 2-year return levels of daily maximum temperature T_X for near future climate (2036-2060 vs. 1990-2014).