
Supplementary information

**Evidence for a vast prograde stellar stream
in the solar vicinity**

In the format provided by the
authors and unedited

Evidence for a Vast Prograde Stellar Stream in the Solar Vicinity

Supplementary Information

Lina Necib,¹ Bryan Ostdiek,² Mariangela Lisanti,³ Timothy Cohen,² Marat Freytsis,^{4,5}
Shea Garrison-Kimmel,⁶ Philip F. Hopkins,⁶ Andrew Wetzel,⁷ and Robyn Sanderson^{8,9}

¹*Walter Burke Institute for Theoretical Physics, California Institute of Technology, Pasadena, CA 91125, USA*

²*Institute of Theoretical Science, Department of Physics, University of Oregon, Eugene, OR 97403, USA*

³*Department of Physics, Princeton University, Princeton, NJ 08544, USA*

⁴*Raymond and Beverly Sackler School of Physics and Astronomy, Tel-Aviv University, Tel-Aviv 69978, Israel*

⁵*School of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540, USA*

⁶*TAPIR, California Institute of Technology, Pasadena, CA 91125, USA*

⁷*Department of Physics, University of California, Davis, CA 95616, USA*

⁸*Department of Physics and Astronomy, University of Pennsylvania, Philadelphia, PA 19104, USA*

⁹*Center for Computational Astrophysics, Flatiron Institute, New York, NY 10010, USA*

This analysis makes use of *emcee* [1] and Extreme Deconvolution [2] for the Gaussian Mixture model.




FIG. 1: **Toomre diagram for Nyx stars.** For reference, the black dashed lines correspond to velocity contours of 100 and 220 km/s in the local frame.




FIG. 2: **Dynamics of the Nyx and Nyx-2 stars.** This figure is similar to Fig. 1, but for stars satisfying a network score cut of $S > 0.85$. In this case, we identify an additional group of stars, labeled Nyx-2, that may be related to the Nyx stream. Details of the clustering algorithm used to find Nyx and Nyx-2 are provided in Paper I. (Left) Plot of the velocities of Nyx and Nyx-2 stars, in the Galactocentric spherical $v_r - v_\phi$ plane. (Middle/Right) Velocity of Nyx and Nyx-2 stars in the $x - y$ and $x - z$ planes, where $x - y$ is the disk plane, and z is perpendicular to the disk. Like Nyx, Nyx-2 is prograde and radial. Its stars move away from the Galactic Center, rather than towards, suggesting that it may be debris from a separate passage of the same satellite galaxy.




FIG. 3: **Network score dependence of the Nyx stars.** This figure is similar to Fig. 1, but for the Nyx stars in the $S > 0.85$ sample. Stars are colored by their neural network score.




FIG. 4: **Skymap of the Nyx stars.** This figure compares Nyx (purple stars) to the footprint of current spectroscopic surveys (gray) using data from [3]. The colorbar indicates the number of stars, which we saturate at 100 for clarity. Nyx has the greatest overlap with the RAVE-ON catalog [4].




FIG. 5: **Chemical abundance of the Nyx stars.** This figure is similar to Fig. 2, but with abundances from RAVE [5] (left) and GALAH [6] (right), rather than RAVE-ON. The error bars are the 1-sigma uncertainties of the surveys.




FIG. 6: **Orbital parameters of the Nyx stars.** The distribution of eccentricities (top left), pericenters (top right), apocenters (bottom left), and maximum vertical distance, $z_{\max}$, from the Galactic plane (bottom right) for the Nyx stars. These parameters were obtained running the orbits of Nyx stars backwards in time over a period of 1 Gyr using the `gala` package [7] and the Milky Way potential of [8].




FIG. 7: **The orbits of the Nyx stars. These are obtained by integrating** back by 1 Gyr using the `gala` package [7] and assuming the Milky Way potential of [8].

- [1] Foreman-Mackey, D., Hogg, D. W., Lang, D. & Goodman, J. emcee: The MCMC Hammer. *Publications of the Astron. Soc. Pacific* **125**, 306 (2013).
- [2] Bovy Jo, Hogg, D. W. & Roweis, S. T. Extreme deconvolution: Inferring complete distribution functions from noisy, heterogeneous and incomplete observations. *Annals of Applied Statistics* **5**, 1657–1677 (2011).
- [3] Sanders, J. L. & Das, P. Isochrone ages for 3 million stars with the second Gaia data release. *Mon. Not. R. Astron. Soc.* **481**, 4093–4110 (2018).
- [4] Casey, A. R. *et al.* The RAVE-on Catalog of Stellar Atmospheric Parameters and Chemical Abundances for Chemo-dynamic Studies in the Gaia Era. *Astrophys. J.* **840**, 59 (2017).
- [5] Kunder, A. *et al.* The Radial Velocity Experiment (RAVE): Fifth Data Release. *Astron. J.* **153**, 75 (2017).
- [6] Buder, S. *et al.* The GALAH Survey: second data release. *Mon. Not. R. Astron. Soc.* **478**, 4513–4552 (2018).
- [7] Price-Whelan, A. M. Gala: A python package for galactic dynamics. *The Journal of Open Source Software* **2** (2017).
- [8] Bovy, J. galpy: A python Library for Galactic Dynamics. *Astrophys. J. Supp.* **216**, 29 (2015).