

In the format provided by the authors and unedited.


The HOIL-1L ligase modulates immune signalling and cell death via monoubiquitination of LUBAC

Yasuhiro Fuseya ^{1,2}, Hiroaki Fujita¹, Minsoo Kim ^{1,3}, Fumiaki Ohtake^{4,5}, Akira Nishide ^{1,3},
Katsuhiro Sasaki ¹, Yasushi Saeki ⁴, Keiji Tanaka⁴, Ryosuke Takahashi ² and Kazuhiro Iwai ¹ 


¹Department of Molecular and Cellular Physiology, Graduate School of Medicine, Kyoto University, Kyoto, Japan. ²Department of Neurology, Graduate School of Medicine, Kyoto University Hospital, Kyoto, Japan. ³The Hakubi Center for Advanced Research, Kyoto University, Kyoto, Japan. ⁴Laboratory of Protein Metabolism, Tokyo Metropolitan Institute of Medical Science, Tokyo, Japan. ⁵Institute for Advanced Life Sciences, Hoshi University, Tokyo, Japan.
✉e-mail: kiwai@mcp.med.kyoto-u.ac.jp

Supplementary Figure1: Gating strategy for flow cytometry analysis


a



b



c



a-c, Single-cell suspensions prepared from the spleen of 8- to 15-week-old mice of the indicated genotype were stained with fluorochrome-conjugated antibodies. Live lymphocytes were gated with forward-scatter (FSC) and side-scatter (SSC), and FSC-area (FSC-A) and FSC-height (FSC-H) were used to discriminate single cells from doublet and multiplet cells. CD19⁺ B cells of splenocytes were analyzed for surface expression of CD38 and PNA, or FAS and PNA (a). TCRβ⁺CD4⁺ T cells of splenocytes were analyzed for surface expression of CD25 (b). Splenocytes were analyzed for surface expression of CD3ε and CD19, and CD19⁻CD3ε⁺ cells were analyzed for surface expression of CD4 and CD8 (c).