

In the format provided by the authors and unedited.

SETD3 is an actin histidine methyltransferase that prevents primary dystocia

Alex W. Wilkinson^{1,9}, Jonathan Diep^{2,9}, Shaobo Dai^{3,9}, Shuo Liu¹, Yaw Shin Ooi², Dan Song⁴, Tie-Mei Li¹, John R. Horton³, Xing Zhang³, Chao Liu⁴, Darshan V. Trivedi⁴, Katherine M. Ruppel⁴, José G. Vilches-Moure⁵, Kerriann M. Casey⁵, Justin Mak⁶, Tina Cowan⁷, Joshua E. Elias⁸, Claude M. Nagamine⁵, James A. Spudich⁴, Xiaodong Cheng^{3,10*}, Jan E. Carette^{2,10*} & Or Gozani^{1,10*}

¹Department of Biology, Stanford University, Stanford, CA, USA. ²Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA, USA. ³Department of Molecular and Cellular Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX, USA. ⁴Department of Biochemistry, Stanford University School of Medicine, Stanford, CA, USA. ⁵Department of Comparative Medicine, Stanford University School of Medicine, Stanford, CA, USA. ⁶Stanford Healthcare, Palo Alto, CA, USA. ⁷Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA. ⁸Department of Chemical and Systems Biology, Stanford University School of Medicine, Stanford, CA, USA. ⁹These authors contributed equally: Alex W. Wilkinson, Jonathan Diep, Shaobo Dai. ¹⁰These authors jointly supervised this work: Xiaodong Cheng, Jan E. Carette, Or Gozani. *e-mail: xcheng5@mdanderson.org; carette@stanford.edu; ogozani@stanford.edu

Figure 1a




Figure 1b




Figure 1d




Figure 1e




Figure 2a





Figure 2c











Figure 3g




Figure 4c




Figure 5g




Figure 5j

