
Supplementary information

**Increased hyaluronan by naked mole-rat
Has2 improves healthspan in mice**

In the format provided by the
authors and unedited

Supplementary Information for

Increased hyaluronan by naked mole-rat HAS2 improves healthspan in mice

Zhihui Zhang¹, Xiao Tian¹, J. Yuyang Lu¹, Kathryn Boit¹, Julia Ablaeva¹, Frances Tolibzoda Zakusilo¹, Stephan Emmrich¹, Denis Firsanov¹, Elena Rydkina¹, Seyed Ali Biashad¹, Quan Lu¹, Alexander Tyshkovskiy^{2, 3}, Vadim N. Gladyshev², Steve Horvath^{4, 5}, Andrei Seluanov^{1, 6, *}, Vera Gorbunova^{1, 6, *}

¹Department of Biology, University of Rochester, Rochester, NY USA

²Division of Genetics, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, 02115, USA

³Belozersky Institute of Physico-Chemical Biology, Moscow State University, Moscow, Russia

⁴Department of Biostatistics, Fielding School of Public Health, University of California Los Angeles, Los Angeles, CA, USA

⁵Department of Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, USA

⁶Department of Medicine, University of Rochester Medical Center, Rochester, NY USA

Contents

Supplementary Discussion	2
Supplementary references	5
Supplementary Table legends	8
Supplementary Table 1. Primer sequences used for quantitative RT-PCR	9

Supplementary Discussion

In the tissues of the naked mole-rat, HMW-HA is abundant and contributes to cancer resistance and possibly longevity of this exceptionally long-lived rodent^{1,2}. Here we demonstrated that this evolutionary adaptation, unique to the naked mole-rat, can be “exported” to other species. The HMW-HA produced by naked mole-rat HAS2 gene conferred cancer resistance and increased lifespan of mice. nmrHAS2 mice accumulated HMW-HA in multiple major organs; these mice were resistant to spontaneous and chemically induced cancer and showed an extended median and maximum lifespan. nmrHAS2 mice also displayed improved healthspan including lower frailty scores and physical performance. This was accompanied by lower methylation age. The transcriptomes of aged nmrHAS mice had more youthful features when compared to the aging signature derived from the Tabula Muris Senis data³. The change conferred by nmrHAS2 was distinct from the transcriptomic changes triggered by known life-extending interventions including rapamycin, acarbose, growth hormone deletion, calorie restrictions and others. Interestingly, the nmrHAS2 altered the mouse transcriptome in a direction consistent with the transcriptomic signature of the long-lived species⁴. We therefore hypothesize that the presence of HMW-HA changes the transcriptome towards that of a longer-lived species (Fig. 4a).

What is the molecular mechanism by which HMW-HA extends mouse lifespan and healthspan? Importantly, due to the high conservation of the HAS2 gene, we do not believe that the naked mole-rat sequence *per se* was critical for the pro-longevity effect of nmrHAS2. Expressing mouse HAS2 gene showed the same protective effects *in vitro*. Rather, the increased production of HMW-HA was important. The most striking difference we observed between the transcriptomes of the aged nmrHAS2 mice and their littermate controls was the downregulation of multiple pathways related to inflammation in most tissues we analyzed. Beneficial effects of HMW-HA have recently been reported for muscle stem cells and adipose tissues where HMW-HA reduced inflammatory signaling and improved glucose homeostasis, respectively^{5,6}. Interestingly, some of the effects were systemic in nature suggesting that even the tissues not displaying elevated HA levels can benefit from elevated HA elsewhere in the body^{5,6}. Additionally, the plasma of aged nmrHAS2 mice also displayed reduced levels of multiple pro-inflammatory cytokines which indicates the systemic anti-inflammatory effect of HMW-HA. Age-associated chronic inflammation, so-called inflammaging, contributes to the pathogenesis of age-related

diseases^{7,8}. It has been reported that individuals which have higher than age-average levels of inflammatory markers are more fragile and likely to be hospitalized⁹.

The lifespan extension we observed in nmrHAS2 mice was significant but modest, while the healthspan improvement was more robust. While we observed strong expression of nmrHAS2 across all tissues, HA accumulation was modest and did not reach the levels observed in the naked mole-rat. This is likely explained by the high activity of hyaluronidases in mouse tissues^{1,10}. In the naked mole-rat high level of HA is achieved by both robust synthesis and very slow degradation¹, yet in our model only the synthesis arm was modified. We hypothesize that if we were able to simultaneously attenuate HA degradation in nmrHAS2 mice we would achieve a greater lifespan extension.

We then discovered two independent pathways by which HMW-HA conferred an anti-inflammatory effect to the mice. First, HMW-HA had a direct immunomodulating effect on the immune cells of nmrHAS2 mice. Monocyte-macrophage lineage cells act as major effector cells in chronic inflammatory processes in aging-related diseases¹¹. Macrophages are often divided into two subgroups: M1 and M2¹². Aging may modulate M1/M2 activation and polarization. Sustained activation of M1 macrophage is linked to tissue dysfunction, while M2 macrophage promotes tissue homeostasis¹³. HMW-HA has been shown to prime macrophages towards the M2 state¹⁴⁻¹⁶. We also observed the same phenotype when we primed the bone marrow-derived macrophages from nmrHAS2 mice with *E. coli* lipopolysaccharide. BMDMs from nmrHAS2 mice showed a much higher expression of HAS2 gene, which may potentially help them to produce more HMW-HA and facilitates the alternative activation. HMW-HA was also shown to play an immunoregulatory role *in vivo* by binding to several immune cell types¹⁷. For instance, HMW-HA promotes the expression of FoxP3 in Treg cells by crosslinking CD44 which helps to maintain immunologic tolerance¹⁸. HMW-HA binds to CD44 and hSiglec-9 receptor which suppresses neutrophil extracellular trap formation and oxidative burst¹⁹. We observed that nmrHAS2 mice produced a lower level of inflammation after the LPS challenge which is consistent with previous reports that intraperitoneal injection of HMW-HA protects mice from LPS-induced sepsis²⁰.

The second, unexpected pathway, by which HMW-HA conferred its anti-inflammatory effect was through improved intestinal health. Aged nmrHAS2 mice were protected from leaky gut, and had healthier microbiomes. One of the major contributors to inflammaging is believed to be age-related deterioration of the intestinal barrier and translocation of gut bacterial products to

the bloodstream triggering an immune response²¹. Intestinal barrier failure is associated with aging-related systemic disorders such as obesity, metabolic disorder, brain dysfunction, and cancer^{22,23}. Reduced thickness of the mucus layers was considered to be the cause of leaky gut²⁴. It has been shown that the colonic mucus layer decreased in aging mice, suggesting an association with bacterial penetration and immune activation^{25,26}. Interestingly, we found that elevated production of hyaluronan increased the number of mucin-producing goblet cells in both small intestine and colon, suggesting old nmrHAS2 are protected from leaky gut due to increased mucin formation. The increase in goblet cells in nmrHAS2 mice was significant, but very mild, and did not resemble a pathological over-proliferation of goblet cells observed in a disease state such as cystic fibrosis²⁷.

Studies in *Drosophila* showed that with aging intestinal stem cell (ISC) lose their stemness, are no longer able to differentiate into functional intestinal cells and undergo hyperproliferation associated with a loss of barrier function²⁸⁻³⁰. While the hyperproliferation of ISCs has not been unequivocally demonstrated in mammals^{28,31}, in mammals aging may be similarly associated with the loss of stemness by the ISCs. Indeed, it was found that the regenerative capacity of ISCs from old mice³¹ and old human²⁸ was diminished *in vitro*. Consistent with these prior reports, we observed that ISCs from aged wild type mice formed fewer intestinal organoids. Remarkably, there was no such decrease observed for nmrHAS2 mice. Interestingly, the decline in the formation of organoids in the wild type mice could be rescued by adding HMW-HA to the culture media, indicating that the presence of HMW-HA promotes stemness of aged ISCs. It was reported that intraperitoneal injection of HA to mice promotes Lgr5+ stem cell proliferation and crypt fission through CD44 and TLR4^{32,33}. Collectively, our results suggest that HMW-HA produced by nmrHAS2 transgene improves the maintenance of ISCs resulting in a healthier gut barrier during aging. Consequently, the healthier gut barrier function inhibits the shift of the gut microbiome towards proinflammatory commensals and reduction of beneficial microbes, slowing down the onset of inflammaging.

The resistance of nmrHAS2 mice to both spontaneous and induced cancer may be driven by both cell-autonomous mechanisms resulting from anti-proliferative signaling of HMW-HA through the CD44 receptor³⁴, and by its systemic anti-inflammatory effect³⁵. Indeed, it was reported that HMW-HA blocks melanoma cell proliferation by signaling through CD44 to promote G1/G0 arrest³⁶. HMW-HA also suppresses the growth of murine astrocytoma cell lines, glioma

and colon carcinoma xenografts³⁷. HMW-HA has also been reported to reduce the migratory and invasive capacity of aggressive cancer cells^{38,39}. On the other hand, chronic inflammation plays an important role in the development of cancer⁷. Many studies have shown that inflammatory cells can promote the occurrence and development of tumors by facilitating cancer cell proliferation, angiogenesis, and tumor invasion⁴⁰⁻⁴². The anti-inflammatory properties of HMW-HA discussed above may potentially reduce the chronic inflammation during aging and prevent cancer initiation.

Additional pro-longevity effects of HMW-HA can be linked to its antioxidant and cytoprotective properties. Indeed, HMW-HA was reported to enhance cellular oxidative stress resistance^{43,44}. Consistent with these observation fibroblasts from nmrHAS2 mice showed resistance to oxidative stress.

Supplementary references

- 1 Tian, X. *et al.* High-molecular-mass hyaluronan mediates the cancer resistance of the naked mole rat. *Nature* **499**, 346-349 (2013).
- 2 Seluanov, A. *et al.* Hypersensitivity to contact inhibition provides a clue to cancer resistance of naked mole-rat. *Proceedings of the National Academy of Sciences* **106**, 19352-19357 (2009).
- 3 Schaum, N. *et al.* Ageing hallmarks exhibit organ-specific temporal signatures. *Nature* **583**, 596-602 (2020).
- 4 Lu, J. Y. *et al.* Comparative transcriptomics reveals circadian and pluripotency networks as two pillars of longevity regulation. *Cell Metabolism* (2022).
- 5 Zhu, Y. *et al.* Adipose tissue hyaluronan production improves systemic glucose homeostasis and primes adipocytes for CL 316,243-stimulated lipolysis. *Nature communications* **12**, 1-15 (2021).
- 6 Nakka, K. *et al.* JMJD3 activated hyaluronan synthesis drives muscle regeneration in an inflammatory environment. *Science* **377**, 666-669 (2022).
- 7 Kennedy, B. K. *et al.* Geroscience: linking aging to chronic disease. *Cell* **159**, 709-713 (2014).
- 8 Franceschi, C. *et al.* Inflamm-aging: an evolutionary perspective on immunosenescence. *Annals of the new York Academy of Sciences* **908**, 244-254 (2000).
- 9 de Gonzalo-Calvo, D. *et al.* Chronic inflammation as predictor of 1-year hospitalization and mortality in elderly population. *European journal of clinical investigation* **42**, 1037-1046 (2012).
- 10 Del Marmol, D. *et al.* Abundance and size of hyaluronan in naked mole-rat tissues and plasma. *Scientific reports* **11**, 1-21 (2021).
- 11 Manabe, I. Chronic inflammation links cardiovascular, metabolic and renal diseases. *Circulation Journal*, 1111041460-1111041460 (2011).

- 12 Martinez, F. O., Gordon, S., Locati, M. & Mantovani, A. Transcriptional profiling of the human monocyte-to-macrophage differentiation and polarization: new molecules and patterns of gene expression. *The Journal of immunology* **177**, 7303-7311 (2006).
- 13 Sica, A., Invernizzi, P. & Mantovani, A. Macrophage plasticity and polarization in liver homeostasis and pathology. *Hepatology* **59**, 2034-2042 (2014).
- 14 Rayahin, J. E., Buhrman, J. S., Zhang, Y., Koh, T. J. & Gemeinhart, R. A. High and low molecular weight hyaluronic acid differentially influence macrophage activation. *ACS biomaterials science & engineering* **1**, 481-493 (2015).
- 15 Bauer, C. *et al.* Chondroprotective effect of high-molecular-weight hyaluronic acid on osteoarthritic chondrocytes in a co-cultivation inflammation model with M1 macrophages. *Journal of Inflammation* **13**, 1-9 (2016).
- 16 Lee, B. M., Park, S. J., Noh, I. & Kim, C.-H. The effects of the molecular weights of hyaluronic acid on the immune responses. *Biomaterials Research* **25**, 1-13 (2021).
- 17 Lee-Sayer, S. S. *et al.* The where, when, how, and why of hyaluronan binding by immune cells. *Frontiers in immunology* **6**, 150 (2015).
- 18 Bollyky, P. L. *et al.* CD44 costimulation promotes FoxP3⁺ regulatory T cell persistence and function via production of IL-2, IL-10, and TGF- β . *The Journal of Immunology* **183**, 2232-2241 (2009).
- 19 Secundino, I. *et al.* Host and pathogen hyaluronan signal through human siglec-9 to suppress neutrophil activation. *Journal of molecular medicine* **94**, 219-233 (2016).
- 20 Muto, J., Yamasaki, K., Taylor, K. R. & Gallo, R. L. Engagement of CD44 by hyaluronan suppresses TLR4 signaling and the septic response to LPS. *Molecular immunology* **47**, 449-456 (2009).
- 21 Thevaranjan, N. *et al.* Age-associated microbial dysbiosis promotes intestinal permeability, systemic inflammation, and macrophage dysfunction. *Cell host & microbe* **21**, 455-466. e454 (2017).
- 22 Chelakkot, C. *et al.* Akkermansia muciniphila-derived extracellular vesicles influence gut permeability through the regulation of tight junctions. *Experimental & molecular medicine* **50**, e450-e450 (2018).
- 23 Biragyn, A. & Ferrucci, L. Gut dysbiosis: a potential link between increased cancer risk in ageing and inflammaging. *The Lancet Oncology* **19**, e295-e304 (2018).
- 24 Cornick, S., Tawiah, A. & Chadee, K. Roles and regulation of the mucus barrier in the gut. *Tissue barriers* **3**, e982426 (2015).
- 25 van Beek, A. A. *et al.* Supplementation with Lactobacillus plantarum WCFS1 prevents decline of mucus barrier in colon of accelerated aging Ercc1- Δ 7 mice. *Frontiers in immunology* **7**, 408 (2016).
- 26 Elderman, M. *et al.* The effect of age on the intestinal mucus thickness, microbiota composition and immunity in relation to sex in mice. *PloS one* **12**, e0184274 (2017).
- 27 Kelly, J. *et al.* Alterations of mucosa-attached microbiome and epithelial cell numbers in the cystic fibrosis small intestine with implications for intestinal disease. *Scientific Reports* **12**, 1-17 (2022).
- 28 Pentimikko, N. *et al.* Notum produced by Paneth cells attenuates regeneration of aged intestinal epithelium. *Nature* **571**, 398-402 (2019).
- 29 Biteau, B., Hochmuth, C. E. & Jasper, H. JNK activity in somatic stem cells causes loss of tissue homeostasis in the aging Drosophila gut. *Cell stem cell* **3**, 442-455 (2008).

- 30 Hochmuth, C. E., Biteau, B., Bohmann, D. & Jasper, H. Redox regulation by Keap1 and Nrf2 controls intestinal stem cell proliferation in *Drosophila*. *Cell stem cell* **8**, 188-199 (2011).
- 31 Nalapareddy, K. *et al.* Canonical Wnt signaling ameliorates aging of intestinal stem cells. *Cell reports* **18**, 2608-2621 (2017).
- 32 Riehl, T. E., Santhanam, S., Foster, L., Ciorba, M. & Stenson, W. F. CD44 and TLR4 mediate hyaluronic acid regulation of Lgr5⁺ stem cell proliferation, crypt fission, and intestinal growth in postnatal and adult mice. *American Journal of Physiology-Gastrointestinal and Liver Physiology* **309**, G874-G887 (2015).
- 33 Riehl, T. E., Alvarado, D., Ee, X., Ciorba, M. A. & Stenson, W. F. Hyaluronic acid promotes Lgr5⁺ stem cell proliferation and crypt fission through TLR4 and PGE2 transactivation of EGFR. *American Journal of Physiology-Gastrointestinal and Liver Physiology* **319**, G63-G73 (2020).
- 34 Kothapalli, D., Flowers, J., Xu, T., Puré, E. & Assoian, R. K. Differential activation of ERK and Rac mediates the proliferative and anti-proliferative effects of hyaluronan and CD44. *Journal of Biological Chemistry* **283**, 31823-31829 (2008).
- 35 Ruppert, S., Hawn, T., Arrigoni, A., Wight, T. & Bollyky, P. Tissue integrity signals communicated by high-molecular weight hyaluronan and the resolution of inflammation. *Immunologic research* **58**, 186-192 (2014).
- 36 Takabe, P. *et al.* Hyaluronan synthase 3 (HAS3) overexpression downregulates MV3 melanoma cell proliferation, migration and adhesion. *Experimental cell research* **337**, 1-15 (2015).
- 37 Mueller, B. M. *et al.* Hyaluronan Inhibits Postchemotherapy Tumor Regrowth in a Colon Carcinoma Xenograft ModelHyaluronan and Postchemotherapy Tumor Regrowth. *Molecular cancer therapeutics* **9**, 3024-3032 (2010).
- 38 Zhao, Y.-f. *et al.* Modulating three-dimensional microenvironment with hyaluronan of different molecular weights alters breast cancer cell invasion behavior. *ACS Applied Materials & Interfaces* **9**, 9327-9338 (2017).
- 39 Enegd, B. *et al.* Overexpression of hyaluronan synthase-2 reduces the tumorigenic potential of glioma cells lacking hyaluronidase activity. *Neurosurgery* **50**, 1311-1318 (2002).
- 40 Coussens, L. M. & Werb, Z. Inflammation and cancer. *Nature* **420**, 860-867 (2002).
- 41 Hanahan, D. & Weinberg, R. A. Hallmarks of cancer: the next generation. *cell* **144**, 646-674 (2011).
- 42 Neagu, M. *et al.* Inflammation: A key process in skin tumorigenesis. *Oncology Letters* **17**, 4068-4084 (2019).
- 43 Onodera, Y., Teramura, T., Takehara, T. & Fukuda, K. Hyaluronic acid regulates a key redox control factor Nrf2 via phosphorylation of Akt in bovine articular chondrocytes. *FEBS Open Bio* **5**, 476-484 (2015).
- 44 Pauloin, T., Dutot, M., Joly, F., Warnet, J.-M. & Rat, P. High molecular weight hyaluronan decreases UVB-induced apoptosis and inflammation in human epithelial corneal cells. *Molecular vision* **15**, 577 (2009).

Supplementary Table legends

Supplementary Table 1. Primer sequences used for quantitative RT-PCR

Supplementary Table 2. CpG islands contribute to younger epigenetics of nmrHAS2 mice

Supplementary Table 3 correlation genes signatures Spearman Padjusted values

Supplementary Table 4. GSEA of enriched functions affected by nmrHAS2 and other interventions

Supplementary Table 5. Fisher functional enrichment of enriched functions affected by nmrHas2 and other interventions

Supplementary Table 1. Primer sequences used for quantitative RT-PCR

Gene	Forward primer	Reverse primer
<i>actb</i>	5'-GGCTGTATTCCCCTCCATCG-3'	5'-CCAGTTGGTAACAATGCCATGT-3'
<i>HAS2</i>	5'-GCCTCATCTGTGGAGATGGT-3'	5'-GAGTTTCTGTACATTCCCAGAGG-3'
<i>Il1b</i>	5'-CCGTGGACCTTCCAGGATGA-3'	5'-GGGAACGTCACACACCAGCA-3'
<i>Il6</i>	5'-AGTTGCCTTCTTGGGACTGA-3'	5'-TCCACGATTTCCCAGAGAAC-3'
<i>tnfa</i>	5'-CATCTTCTCAAAATTCGAGTGACAA-3'	5'-TGGGAGTAGACAAGGTACAACCC-3'
<i>arg1</i>	5'-GCTCAGGTGAATCGGCCTTTT-3'	5'-TGGCTTGCGAGACGTAGAC-3'
<i>Il12b</i>	5'-AGACCCTGCCCATTGAACTG-3'	5'-GAAGCTGGTGCTGTAGTTCTCATATT-3'
<i>nos2</i>	5'-TTCACCCAGTTGTGCATCGACCTA-3'	5'-TCCATGGTCACCTCCAACACAAGA-3'
<i>Il10</i>	5'-GAGAGCTGCAGGGCCCTTTGC-3'	5'-CTCCCTGGTTTCTCTTCCCAAGACC-3'
<i>Hyal1</i>	5'-CATGCCTGAACCTGACTTCT-3'	5'-GTAGCAGTCAGGGAAGCCATA-3'
<i>Hyal2</i>	5'-CACCTGCCCATGCTGAAGGA-3'	5'-TCAGGAAAGAGGTAGAAGCC-3'