

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see [Authors & Referees](#) and the [Editorial Policy Checklist](#).

Statistical parameters

When statistical analyses are reported, confirm that the following items are present in the relevant location (e.g. figure legend, table legend, main text, or Methods section).

n/a Confirmed

- ☐ ☒ The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- ☐ ☒ An indication of whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- ☐ ☒ The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- ☐ ☒ A description of all covariates tested
- ☐ ☒ A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- ☐ ☒ A full description of the statistics including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- ☐ ☒ For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- ☒ ☐ For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- ☒ ☐ For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- ☐ ☒ Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated
- ☐ ☒ Clearly defined error bars
State explicitly what error bars represent (e.g. SD, SE, CI)

Our web collection on [statistics for biologists](#) may be useful.

Software and code

Policy information about [availability of computer code](#)

Data collection

RNA seq, Microarray and qRT-PCR data were collected using vendor's software on Illumina HiSeq 2000, Agilent Whole Human Oligo Microarray, Applied Biosystems 7900HT Real-Time PCR platforms respectively. Microscopy images were acquired using Metamorph.

Data analysis

Software for transcriptome meta-assembly and lncRNAs discovery is available at <https://tacorna.github.io/>. Gene signatures were obtained using GSEA software. Image analysis was performed using custom-written macros in Image J and can be shared upon request. Statistical analysis was performed using Graphpad-Prism 6.0 and R.

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors/reviewers upon request. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

RNA-seq and microarray datasets generated from this study have been deposited into Gene Expression Omnibus, with accession number: GSE110905. Other data supporting the finding of this study are included in the Supplementary Information files. Software for transcriptome meta-assembly and lncRNAs discovery is available at <https://tacorna.github.io/>. We have no restrictions on data availability and data can be shared upon request.

Field-specific reporting

Please select the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☒ Life sciences ☐ Behavioural & social sciences

For a reference copy of the document with all sections, see nature.com/authors/policies/ReportingSummary-flat.pdf

Life sciences

Study design

All studies must disclose on these points even when the disclosure is negative.

Sample size	For in vivo experiments, power analysis (GPOWER software) performed for each tumor type tested to date indicates this animal number yields a statistical power >90% for detection of tumor size reduction of 40%. Sample sizes were not pre-determined for all other assays.
Data exclusions	No data were excluded.
Replication	All experiments were carried out at least 3 independent times for statistical reproducibility, unless otherwise stated, as represented by p-values.
Randomization	For in vivo experiments, animals were randomized. Randomization was not performed for all other assays.
Blinding	For in vivo experiments, tumor size was measured twice per week using a digital caliper by a researcher blinded to the study design. Blinding was not performed for all other assays.

Materials & experimental systems

Policy information about [availability of materials](#)

n/a	Involved in the study
☒	☐ Unique materials
☐	☒ Antibodies
☐	☒ Eukaryotic cell lines
☐	☒ Research animals
☒	☐ Human research participants

Antibodies

Antibodies used	Primary antibodies used in this study were: Androgen Receptor (1:1000 dilution, Millipore, #06-680, rabbit), GAPDH (1:5000 dilution, Cell Signaling, #3683, rabbit), PSA (1:5000 dilution, Dako, #A0562, rabbit), FOXA1 (Thermo Fisher Cat# PA5-27157) NKX3.1 (CST Cat# 837005) and cleaved PARP (1:1000 dilution, Cell Signaling, #9542, rabbit).
Validation	All antibodies were validated by the vendors. Androgen receptor and PSA antibodies were also validated by siRNA treatment and androgen signaling assays respectively.

Eukaryotic cell lines

Policy information about [cell lines](#)

Cell line source(s)	All cell lines were purchased from ATCC.
Authentication	All cell lines were genotyped by STR profiling in house based on ATCC markers.

Mycoplasma contamination	All cells were tested for mycoplasma every two weeks.
Commonly misidentified lines (See ICLAC register)	Study does not include misidentified lines.

Research animals

Policy information about [studies involving animals](#); [ARRIVE guidelines](#) recommended for reporting animal research

Animals/animal-derived materials	Male athymic nude mice were used in our in vivo studies.
----------------------------------	--

Method-specific reporting

n/a	Involved in the study
☒	☐ ChIP-seq
☒	☐ Flow cytometry
☒	☐ Magnetic resonance imaging