

In the format provided by the authors and unedited.

Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding

In the format provided by the authors and unedited

In the format provided by the authors and unedited.

Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding

Yi Xu^{1,11}, Xufang Li^{1,11}, Bing Zhu^{2,11}, Huiying Liang^{3,4,11}, Chunxiao Fang¹, Yu Gong¹, Qiaozhi Guo⁵, Xin Sun⁵, Danyang Zhao⁵, Jun Shen¹, Huayan Zhang^{1,4,6}, Hongsheng Liu⁷, Huimin Xia^{3,4,8}✉, Jinling Tang^{3,4}✉, Kang Zhang^{8,9,10}✉ and Sitang Gong^{1,4}✉

¹Department of Pediatric, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, China. ²Department of Center Laboratory, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, China. ³Clinical Data Center, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, China. ⁴Guangdong Provincial Children's Medical Research Center, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, China. ⁵Department of Medical Administration, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, China. ⁶Division of Neonatology and Department of Pediatrics, Children's Hospital of Philadelphia and University of Pennsylvania, Philadelphia, PA, USA. ⁷Department of Radiology, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, China. ⁸Guangdong Provincial Key Laboratory of Research in Structural Birth Defect Disease, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, China. ⁹Guangzhou Regenerative Medicine and Health Guangdong Laboratory, Guangzhou, China. ¹⁰Center for Biomedicine and Innovations, Faculty of Medicine, Macau University of Science and Technology, Macau, China. ¹¹These authors contributed equally: Yi Xu, Xufang Li, Bing Zhu, Huiying Liang. ✉ e-mail: huiminxia@hotmail.com; jltang@cuhk.edu.hk; kang.zhang@gmail.com; sitangg@126.com

Supplementary Table 1. Cytokine levels of n=10 independent pediatric patients confirmed with SARS CoV-2 infection at hospital admission

Characteristic	Case 1	Case 2	Case 3	Case 4	Case 5	Case 6	Case 7	Case 8	Case 9	Case 10
IgG (g/L; normal range 5-10.6)	9.25	11.90	11	16.00	6.02	8.28	18.1	15.1	3.73	NA
IgA (g/L; normal range 0.34-1.38)	1.34	2.49	0.86	3.78	0.37	0.6	1.82	2.61	0.07	NA
IgM (g/L; normal range 0.44-1.44)	0.95	1.33	1.19	0.87	0.79	0.53	1.81	1.57	0.36	NA
C3 (g/L; normal range 0.8-1.5)	1.06	0.96	0.89	0.72	0.94	0.99	1.37	1.38	0.66	NA
C4 (g/L; normal range 0.13-0.43)	0.3	0.22	0.27	0.15	0.26	0.23	0.32	0.25	0.07	NA
IgE (g/L; normal range 0-100)	638	55	144	411	41	221	18	50	15	NA
TNF- β (ng/L; normal range 0-3.49)	0.73	1.21	0.72	1.72	1.06	0.92	2.02	1.43	1.91	NA
IL-12p70 (ng/L; normal range 0-6.9)	6.12	7.06	2.76	5.90	5.31	3.01	2.94	2.83	3.54	NA
IL-1 β (ng/L; normal range 0-3.12)	0.67	0.90	0.15	2.69	0.39	0.63	0.01	1.95	0.01	NA
IL-10 (ng/L; normal range 0-8.14)	3.9	4.10	2.99	6.78	13.31	4.2	2.57	2.90	4.57	NA
IL-6 (ng/L; normal range 0-8.88)	8.62	14.25	60.09	14.40	8.47	8.93	4.46	16.58	2.23	NA
TNF- α (ng/L; normal range 0-5.35)	4.34	5.26	2.99	9.04	13.90	4.34	3.46	4.29	3.76	NA
IL-2 (ng/L; normal range 0-5.03)	8.73	6.19	4.85	10.49	72.91	6.28	1.59	2.39	3.73	NA
IFN- γ (ng/L; normal range 0-6.56)	4.09	4.82	2.72	6.65	11.82	2.8	2.82	5.56	5.76	NA
IL-17F (ng/L; normal range 0-4.66)	6.42	7.29	4.91	11.95	8.90	4.73	3.39	5.28	4.32	NA
IL-8 (ng/L; normal range 0-15.71)	4.42	3.96	12.21	9.00	6.20	21.29	4.67	3.68	5.18	NA
IL-4 (ng/L; normal range 0-4.62)	2.08	1.58	1.58	5.72	3.41	1.83	1.62	2.50	1.83	NA
IL-5 (ng/L; normal range 0-3.73)	1.38	1.03	0.49	2.36	0.99	1.17	1.31	2.19	1.40	NA
IL-17A (ng/L; normal range 0-3.71)	3.31	3.12	2.22	3.96	2.86	2.79	2.37	2.87	3.96	NA
IL-22 (ng/L; normal range 0-2.61)	1.78	5.81	10.22	9.71	4.86	2.11	0.75	3.98	2.18	NA

IL= Interleukin