

ROLE OF NEW IMMUNOPHENOTYPIC MARKERS ON PROGNOSTIC AND OVERALL SURVIVAL OF ACUTE MYELOID LEUKEMIA: A SYSTEMATIC REVIEW AND META-ANALYSIS

Costa AFO¹, Menezes DL¹, Pinheiro LHS¹, Sandes AF², Nunes MAP³, Lyra Junior DP¹,
Schimieguel DM¹

¹Department of Pharmacy, Laboratory of Hematology, Federal University of Sergipe,
Aracaju, Sergipe, Brazil.


²Fleury Group, Hematology Division, São Paulo, São Paulo, Brazil.

³Department of Medicine, Federal University of Sergipe, Aracaju, Sergipe, Brazil.

*Corresponding author:


amandafernandes@hotmail.com (AF)

Supplementary Figure S1. Forest plot with relative risks and confidence intervals of survival in 10 months.




Relative risks and confidence intervals of survival at 10 months associated to the non-detection / detection of the immunophenotypic markers in each study and its meta-analytical measurements before the withdraw of the two studies found to be responsible for the asymmetry.

Supplementary Figure S2. Funnel plot of studies on survival in 10 months.




Publication bias potential of 10-month survival associated with the non-detection / detection of the immunophenotypic markers in each study and its meta-analytical measurements before the withdraw of the two studies found to be responsible for the asymmetry.

Supplementary Figure S3. Funnel plot of studies on survival in 10 months



Publication bias potential of 10-month survival associated with the non-detection / detection of the immunophenotypic markers in each study and its meta-analytical measurements after the withdraw of the two studies found to be responsible for the asymmetry.

Supplementary Figure S4. Funnel plot of studies on survival in 10 months.



Publication bias potential of 20-month survival associated with the non-detection / detection of the immunophenotypic markers in each study and its meta-analytical measurements.