

The adjuvant value of *Andrographis paniculata* in metastatic esophageal cancer treatment – from preclinical perspectives

Lin Li¹, Grace Gar-Lee Yue², Julia Kin-Ming Lee², Eric Chun-Wai Wong², Kwok-Pui Fung^{2,3}, Jun Yu⁴, Clara Bik-San Lau^{2,*}, Philip Wai-Yan Chiu^{1,*}

¹ Department of Surgery; ² Institute of Chinese Medicine and State Key Laboratory of Phytochemistry and Plant Resources in West China (CUHK); ³ School of Biomedical Sciences; ⁴ Department of Medicine and Therapeutics and State Key Laboratory of Digestive Disease, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong.

Supporting Information

Table S1. Sequence of specific PCR primers of different target genes

<i>Genes</i>	Forward Primer	Reverse Primer
<i>TM4SF3</i>	ATAGATATCGACAAGCCTGTAACGAA	GATCTCGAGGTTCCCGATCTGG
<i>MMP9</i>	Hs_MMP9_1_SD QuantiTect Primer Assay (QIAGEN)	
<i>HER2</i>	GAGCCGCGAGCACCCAAGT	TCCATTGTCTAGCACGGCCA
<i>CXCR4</i>	GGCAGAGGAGTTAGCCAAGAT	CTCCTCCCCATCTTTTCCCAT
<i>GAPDH</i>	CGAGATCCCTCCAAAATCAA	GGTGCTAAGCAGTTGGTGGT