

**Identification and activation of TLR4-mediated signalling pathways by
alginate-derived guluronate oligosaccharide in RAW264.7 macrophages**

Weishan Fang ^{1,*}, Decheng Bi ^{1,*}, Ruijin Zheng ^{2,*}, Nan Cai ¹, Hong Xu ¹, Rui Zhou ¹,
Jun Lu ^{1,3}, Min Wan ⁴, Xu Xu ¹

¹ College of Life Sciences and Oceanography, Shenzhen Key Laboratory of Marine Bioresources and Ecology, Shenzhen University, Shenzhen 518060, PR China.

² Department of Environmental & Community Medicine, Rutgers University-Robert Wood Johnson Medical School, Piscataway, NJ 08854, USA.

³ School of Science and School of Interprofessional Health Studies, Faculty of Health and Environmental Sciences, and Institute of Biomedical Technology, Auckland University of Technology, Auckland 1142, New Zealand.

⁴ Division of Physiological Chemistry 2, Department of Medical Biochemistry and Biophysics, Karolinska Institute, Stockholm 17177, Sweden.

Correspondence and requests for materials should be addressed to X. X. (email: xuxu@szu.edu.cn)

* These authors contributed equally to this work.

Figure 2C




Figure 2D-E




Figure 3A




Figure 3B




Figure 3C




Figure 4A




Figure 4B




Figure 5A-F

