

Supplementary Information

Generation of *Myostatin* Gene-Edited Channel Catfish (*Ictalurus punctatus*) via Zygote Injection of CRISPR/Cas9 System

Karim Khalil^{1, 2,*}, Medhat Elayat², Elsayed Khalifa ², Samer Daghash², Ahmed Elaswad^{1, 4,*}, Michael Miller³, Hisham Abdelrahman^{1,5}, Zhi Ye¹, Ramjie Odin¹, David Drescher¹, Khoi Vo¹, Kamal Gosh¹, William Bugg¹, Dalton Robinson¹ & Rex Dunham^{1,*}

¹School of Fisheries, Aquaculture and Aquatic Sciences, Auburn University, Auburn, AL, 36849, USA.

²Anatomy and Embryology Department, Faculty of Veterinary Medicine, Cairo University, Giza 12211, Egypt.


³Harrison School of Pharmacy, Auburn University, Auburn, AL, 36849, USA.

⁴Department of Animal Wealth Development, Faculty of Veterinary Medicine, Suez Canal University, Ismailia 41522, Egypt.

⁵Department of Veterinary Hygiene and Management, Faculty of Veterinary Medicine, Cairo University, Giza 12211, Egypt.

*Correspondence and requests for materials should be addressed to Karim Khalil, Karim.Khalil@vet.cu.edu.eg / Karim11ane@gmail.com, Ahmed Elaswad, ahe0001@auburn.edu, Rex Dunham, dunhara@auburn.edu, School of Fisheries, Aquaculture and Aquatic Sciences, Auburn University, 203 Swingle Hall, Auburn, Alabama 36849, United States. Tel: +1 3348444786; Fax: +1 3348449208.

Figure S1



Supplementary Figure S1 Original images of gels.

Results of Surveyor mutation detection in Fig. 3A (A, B) were shown. Red box represent cropped area.