

CitH3: a reliable blood biomarker for diagnosis and treatment of endotoxic shock

Baihong Pan^{1,2}, Hasan B. Alam¹, Wei Chong³, Jim Mobley⁴, Baoling Liu¹, Qiufang Deng^{1,2},
Yinjian Liang^{1,3}, Yanming Wang⁵, Eric Chen¹, Tianbing Wang⁶, Muneesh Tewari⁷, and
Yongqing Li^{1*}

¹ Department of Surgery, University of Michigan Hospital, Ann Arbor, MI, USA

² Department of General Surgery, Xiangya Hospital, Changsha, Hunan, China

³ The First Hospital, China Medical University, Shenyang, China

⁴ Cayman Chemical, Ann Arbor, MI, 48108, USA

⁵ Department of Biochemistry and Molecular Biology, Penn State University, University Park, PA, US.

⁶ Department of Trauma and Orthopedic Surgery, Peking University People's Hospital, Beijing, China

⁷ Department of Internal Medicine, University of Michigan, Ann Arbor, MI, USA

*Corresponding author:

Yongqing Li, MD, PhD

Email: yqli@med.umich.edu

Department of Surgery


University of Michigan Medical School

NCRC, Room 363N, Building 26

Plymouth Road, Ann Arbor, MI, USA

LPS model


		<u>0.5h</u>		<u>3h</u>		<u>12h</u>		<u>24h</u>	
LPS	-	+	+	+	+	+	+	+	+
YW3-56	-	-	+	-	+	-	+	-	+



17KD

HS model

	<u>Hours</u>				
Sham	0.5	3	12	24	CTL(+)



17KD

Supplementary Figure 1. Effect of different treatments (LPS, LPS+YW3-56, or HS) on blood levels of CitH3 in mouse models of LPSS and HS. Full-length western blots of circulating CitH3 which corresponds to cropped blots presented in Figure 2 (A and C). See the figure legend of Figure 2 for detail.