

Location of the Central Retinal Vessel Trunk in the Lamina and Prelamina Tissue of Healthy and Glaucomatous Eyes

Authors: Bo Wang^{1,2*}, Katie A Lucy^{3*}; Joel S Schuman³; Hiroshi Ishikawa³; Richard A Bilonick^{1,4}; Ian A Sigal^{1,2}; Larry Kagemann⁵; Chen Lu⁶; James G Fujimoto⁶; Gadi Wollstein³

Affiliations: ¹UPMC Eye Center, Eye and Ear Institute, Ophthalmology and Visual Science Research Center, Department of Ophthalmology, University of Pittsburgh School of Medicine, Pittsburgh, PA;

²Department of Bioengineering, Swanson School of Engineering, University of Pittsburgh, Pittsburgh, PA;

³NYU Langone Eye Center, New York University School of Medicine, New York, NY;

⁴Department of Biostatistics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA;

⁵Center for Devices and Radiological Health, Food and Drug Administration, Silver Spring, MD;

⁶Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology, Cambridge, MA.

*These authors contributed equally to this work.

Supplemental Video 1

Example of CRVT tracing in a subject with thin prelaminae tissue. An OCT B-scan (left panel) shows the location (yellow line) of the C-mode (middle panel). The location of the CRVT is traced (right panel) with a yellow circle.

Supplemental Video 2

Example of CRVT tracing in a subject with thick prelaminae tissue. An OCT B-scan (left panel) shows the location (yellow line) of the C-mode (middle panel). The location of the CRVT is traced (right panel) with a yellow circle.