

Comparison of prognostic, clinical, and renal histopathological characteristics of overlapping idiopathic membranous nephropathy and IgA nephropathy versus idiopathic membranous nephropathy

Xinxin Chen¹, Yu Chen², Keqing Shi³, Yinqiu Lv¹, Huan Tong⁴, Guangju Zhao⁴,
Chaosheng Chen¹, Bo Chen¹, Duo Li¹, Zhongqiu Lu⁴ *

Supporting information:

Table S1. The association between different treatments and poor prognosis. Factors predicting prognosis in univariate Cox regression models.

Factor	Univariate Cox Model	
	Hazard ratio (95%CI)	<i>P</i> value
Different treatment	0.698 (0.472-1.033)	0.072

Poor prognosis was defined as an occurrence of a permanent 50% reduction in eGFR, ESRD, or all-cause mortality. $P < 0.05$ was considered statistically significant. CI: confidence interval.

Table S2. The association between different treatments and good prognosis. Factors predicting prognosis in univariate Cox regression models.

Factors	Univariate Cox Model	
	Hazard ratio (95%CI)	<i>P</i> value
Different treatment	0.927 (0.811-1.061)	0.270

Good prognosis was defined as an occurrence of PR or CR. $P < 0.05$ was considered statistically significant. CI: confidence interval.