

Supplementary information

Two NAD-linked redox shuttles maintain the peroxisomal redox balance in *Saccharomyces cerevisiae*

Nadal A. Al-Saryi^{1,2}, Murtakab Y Al-Hejjaj¹, Carlo W. T. van Roermund³, Georgia E. Hulmes¹, Lakhan Ekal¹, Chantell Payton⁴, Ronald J. A. Wanders³ and Ewald H. Hettema¹

1 Department of Molecular Biology, University of Sheffield, UK

2 Current address, Department of Biology, College of Science, Al Mustansiriyah University, Bagdad, Iraq

3 Laboratory Genetic Metabolic Diseases, Department of Clinical Chemistry, Academic Medical Center, Amsterdam, The Netherlands

4 School of Life Sciences, University of Lincoln, UK

Corresponding author:

E.H. Hettema

Email: e.hettema@sheffield.ac.uk

Supplementary table 1. Yeast strains used in this study

Strain	Source
BY4741 MATa <i>his3-1 leu2-0 met15-0 ura3-0</i>	Euroscarf
BY4742 MATa <i>his3-1 leu2-0 lys2-0 ura3-0</i>	Euroscarf
BY4741 <i>fox1::kan^r</i>	Euroscarf
BY4741 <i>fox3::kan^r</i>	Euroscarf
BY4741 <i>gpd1::kan^r</i>	Euroscarf
BY4741 <i>mdh3::SchizHIS5</i>	This study
BY4741 <i>pex3::kan^r</i>	Euroscarf
BY4741 <i>gpd1::kan^r, mdh3::SchizHIS5</i>	This study
BY4741 <i>pex3::hph^r, gpd1::kan^r, mdh3::SchizHIS5</i>	This study
BY4741 <i>pex5::hph^r, gpd1::kan^r, mdh3::SchizHIS5</i>	This study
BY4741 <i>pex7::hph^r, gpd1::kan^r, mdh3::SchizHIS5</i>	This study
BY4741 <i>gpd1::kan^r, mdh3::SchizHIS5 LYS1-mCherry::hph^r</i>	This study

Supplementary figure 1

Full sized western blots from which the compilation in Fig. 2 was made

