

Polydatin inhibits mast cell-mediated allergic inflammation by targeting PI3K/Akt, MAPK, NF-κB and Nrf2/HO-1 pathways

Jing Ye^{1,#}, Hongmei Piao^{2,#}, Jingzhi Jiang¹, Guangyu Jin², Mingyu Zheng³, Jinshi Yang³, Xiang Jin³, Tianyi Sun³, Yun Ho Choi⁴, Liangchang Li^{1,*}, Guanghai Yan^{1,*}

¹Department of Anatomy and Histology and Embryology, Yanbian University Medical College, Yanji 133002, P.R. China;

²Department of Respiratory Medicine, Yanbian University Hospital, Yanji, P.R. China;

³College of Pharmacy, Yanbian University, Yanji 133002, P.R. China;

⁴Department of Anatomy, Medical School, Institute for Medical Sciences, Chonbuk National University, Jeonju, Jeonbuk 561-756, Republic of Korea

#These authors contributed equally to this work.

*Corresponding authors:

Guanghai Yan: Department of Anatomy, Histology and Embryology, Yanbian University Medical College, No. 977 Gongyuan Road, Yanji 133002, Jilin Province, P.R. China

Tel.: 86-433-2435137

Fax: 86-433-2435136

Email: ghyan@ybu.edu.cn & ghyan2015@sina.com

Liangchang Li: Department of Anatomy, Histology and Embryology, Yanbian University Medical College, No. 977 Gongyuan Road, Yanji 133002, Jilin Province, P.R. China

Tel.: 86-433-2435137

Fax: 86-433-2435136

Email: lcli@ybu.edu.cn

Figure 3A

Fig. 3A




Figure 4

Fig. 4




Figure 5

Fig. 5




Figure 6

Fig. 6





Figure 7

Fig. 7
A



B

