

Supplementary material

Potential Application of ^1H NMR for Routine Serum Lipidome Analysis – Evaluation of Effects of Bariatric Surgery

Adriana Mika^{1,2,*}, Zbigniew Kaczynski³, Piotr Stepnowski¹, Maciej Kaczor⁴, Monika Proczko-Stepaniak⁴, Lukasz Kaska⁴, Tomasz Sledzinski²

¹ Department of Environmental Analysis, Faculty of Chemistry, University of Gdansk, Wita Stwosza 63, 80-308 Gdansk, Poland

² Department of Pharmaceutical Biochemistry, Medical University of Gdansk, Debinki 1, 80-211 Gdansk, Poland

³ Faculty of Chemistry, University of Gdansk, Wita Stwosza 63, 80-308 Gdansk, Poland

⁴ Department of General, Endocrine and Transplant Surgery, Medical University of Gdansk, Smoluchowskiego 17, 80-214 Gdansk, Poland

Corresponding author:

* Dr. Adriana Mika; Department of Environmental Analysis, Faculty of Chemistry, University of Gdansk, Wita Stwosza 63, Gdansk, 80-308; Poland; tel./fax: +48585230810,

Email: adrianamika@tlen.pl




Figure S2. The ^1H - ^1H TOCSY spectrum of MO patients' serum determining the position of TG and PL protons. TG - violet lines; PL - red lines.