

Supplementary Results to Accompany

Brain Structural Bases of Tendency to Forgive: evidence from a young adults sample using voxel-based morphometry

Haijiang Li^{1*}, Qunlin Chen^{2,3*}, Jiamei Lu¹, Jiang Qiu^{2,3}

¹Department of Psychology, Shanghai Normal University, Shanghai 200234, China, ²Key Laboratory of Cognition and Personality (SWU), Ministry of Education, Chongqing 400715, China, ³Faculty of Psychology, Southwest University, Chongqing 400715, China.

Correspondence authors:

Jiamei Lu, Prof. (lujiameivip@126.com)

Department of Psychology, Shanghai Normal University
No.100 Guilin Rd. Xuhui district, Shanghai, China, 200234

Tel: +86 21 6432 3257

Table S1 Brain regions with significant associations between TTF and cortical thickness.

Brain regions	Side	MNI coordination			NVtxs	Peak
		x	y	z		T-Value
<i>Positive correlations</i>						
Superior Frontal Gyrus	R	20	-4	61	203	3.25
IFG	L	-49	10	17	249	3.41
<i>Negative correlation</i>						
Superior Frontal Gyrus	R	19	28	49	134	2.99
IPL	R	45	-49	41	149	2.89
IFG	R	44	27	21	166	2.84
Insular cortex	R	39	-7	1	121	2.48

Statistical threshold, $p < 0.01$, uncorrected. TTF, tendency to forgive; IFG, inferior frontal gyrus; Inferior parietal lobule, IPL; NVtxs, number of vertex.

Figure S1 Association between TTF scores and cortical thickness.

