

Seasonal Expression of Oxytocin and Oxytocin Receptor in the Scented Gland of Male Muskrat (*Ondatra zibethicus*)

Fengwei Zhang¹, Qian Liu¹, Ziyi Wang¹, Wenqian Xie¹, Xia Sheng², Haolin Zhang¹, Zhengrong Yuan¹, Yingying Han^{1*}, Qiang Weng^{1*}.

¹*College of Biological Sciences and Technology, Beijing Forestry University, 100083 Beijing, China.*

²*Department of Biosciences, University of Oslo, 0316 Oslo, Norway.*

*Correspondence:

Yingying Han, Ph.D., Laboratory of Animal Physiology, College of Biological Sciences and Technology, Beijing Forestry University, Beijing 100083, China. Fax: +86 10-6233-6399; E-mail: thinkinghy@bjfu.edu.cn

Qiang Weng, Ph.D., Laboratory of Animal Physiology, College of Biological Sciences and Technology, Beijing Forestry University, Beijing 100083, China. Fax: +86 10-6233-6399; E-mail: qiangweng@bjfu.edu.cn

Supplementary Figure S1




Figure S1: Full-length blots of Figure 4. **(a)** corresponds to Figure 4a; **(b)** corresponds to Figure 4b; **(c)** corresponds to Figure 4c; **(d)** corresponds to Figure 4d. B, breeding season; NB, non-breeding season; NC, negative control.

Supplementary Figure S2




Figure S2: (a-c) Three different DNA markers. (a) corresponds to *Oxt*, *Oxtr* and *Mapk3*; (b) corresponds to *Mapk1* and *Actb*; (c) corresponds to *Fos*. (d) Original agarose gels of *Oxt*, *Oxtr*, *Mapk3*, *Mapk1*, *Fos* and *Actb* mRNA, respectively. White rectangles indicate the lanes assembled in the Figure 5. B, breeding season; NB, non-breeding season.