

Comparative toxicity of five dispersants to coral larvae

Negri, A.P.^{1*}, Luter, H.M.¹, Fisher, R.¹, Brinkman, D.L.¹, Irving, P.²

¹Australian Institute of Marine Science, Townsville, QLD, and Perth, WA, Australia

²Australian Maritime Safety Authority, Canberra, ACT, Australia

* Corresponding author

email: a.negri@aims.gov.au

Table S-1. Water quality parameters measured at the end of the longest exposure period (24 h).

Dissolved oxygen (DO, mg l⁻¹) was measured for each treatment, while pH and salinity were measured for the seawater control and 40 mg l⁻¹ treatments.

Treatment	Concentration (mg dispersant l ⁻¹)								
	0	0.1	1.25	2.5	5	10	20	40	80
<i>DO</i>									
Seawater	7.6	7.5	7.7	7.2	7.2	7.6	7.4	7.4	7.5
Corexit EC9500A	7.6	7.5	7.6	7.3	7.4	7.5	7.4	7.5	7.5
Slickgone LTSW	7.5	7.5	7.3	7.3	7.2	7.4	7.3	7.4	7.2
Slickgone NS	7.5	7.4	7.1	7.2	7.3	7.6	7.4	7.2	7.1
Ardrox	7.6	7.3	7.4	7.5	7.4	7.4	7.5	7.2	7.1
Finisol	7.6	7.4	7.6	7.6	7.4	7.5	7.5	7.4	7.2
SDS	7.4	7.4	7.5	7.4	7.3	7.4	7.2	7.1	7.1
<i>pH</i>									
Control	8.11							8.15	
Corexit EC9500A	8.17							8.16	
Slickgone LTSW	8.09							8.07	
Slickgone NS	8.11							8.12	
Ardrox	8.08							8.13	
Finisol	8.14							8.14	
SDS	8.06							8.10	
<i>salinity</i>									
Control	34							33	
Corexit EC9500A	34							34	
Slickgone LTSW	35							35	
Slickgone NS	33							34	
Ardrox	34							34	
Finisol	34							34	
SDS	33							33	