

Gene-based analysis of genes related to neurotrophic pathway suggests association of *BDNF* and *VEGFA* with antidepressant treatment-response in depressed patients

Chung-Feng Kao¹, Yu-Li Liu², Younger W-Y Yu³, Albert C. Yang^{4,5,6,7}, Eugene Lin^{8,9}, Po-Hsiu Kuo^{10*} & Shih-Jen Tsai^{4,5,7*}

¹ Department of Agronomy, National Chung Hsing University, Taichung, Taiwan

² Center for Neuropsychiatric Research, National Health Research Institutes, Miaoli County, Taiwan

³ Yu's Psychiatric Clinic, Kaohsiung, Taiwan

⁴ Department of Psychiatry, Taipei Veterans General Hospital, Taipei, Taiwan

⁵ Division of Psychiatry, National Yang-Ming University, Taipei, Taiwan

⁶ Division of Interdisciplinary Medicine and Biotechnology, Beth Israel Deaconess Medical Center/Harvard Medical School, Boston, MA 02215, USA

⁷ Institute of Brain Science, National Yang-Ming University, Taipei, Taiwan

⁸ Graduate Institute of Biomedical Sciences, China Medical University, Taichung, Taiwan

⁹ Department of Electrical Engineering, University of Washington, Seattle, WA 98195, USA

¹⁰ Department of Public Health, Institute of Epidemiology and Preventive Medicine, National Taiwan University, Taipei, Taiwan

*** Corresponding author:**

E-mail: phkuo@ntu.edu.tw (P.-H.K.)

or

E-mail: tsai610913@gmail.com (S.-J.T.)

Short Title: Gene-based analysis of antidepressant effect

Figure S1




Figure S2




Figure S3

Mahattan plot of selected candidate-genes for Response (%ΔHRSD, binary)



Figure S4

Mahattan plot of selected candidate-genes for Response (%ΔHRSD, binary)




Figure S5

Mahattan plot of selected candidate-genes for Stem-depressed (itemwise)




Figure S6

Mahattan plot of selected candidate-genes for Stem-depressed (itemwise)

