

The neurotrophic tyrosine kinase receptor TrkA and its ligand NGF are increased in squamous cell carcinomas of the lung.

Fangfang Gao^{1,2}, Nathan Griffin^{1,2}, Sam Faulkner^{1,2}, Christopher W. Rowe^{2,3}, Lily Williams¹, Severine Roselli^{1,2}, Rick F. Thorne², Aysha Ferdoushi^{1,2}, Phillip Jobling^{1,2}, Marjorie M. Walker^{2,3}, Hubert Hondermarck^{1,2,*}



¹School of Biomedical Sciences & Pharmacy, Faculty of Health and Medicine, University of Newcastle, Callaghan NSW 2308, Australia.

²Hunter Medical Research Institute, University of Newcastle, New Lambton NSW 2305, Australia.

³School of Public Health & Medicine, Faculty of Health and Medicine, University of Newcastle, Callaghan NSW 2308, Australia.

Corresponding author: Hubert Hondermarck, School of Biomedical Sciences & Pharmacy, Life Sciences Building (LS3-35), University of Newcastle, Callaghan NSW 2308, Australia. Tel: +61 2492 18830. Email: hubert.hondermarck@newcastle.edu.au

Supplementary Figure 1: Negative controls for immunohistochemistry. A. Rabbit isotype control antibody was used in immunohistochemistry for NGF (ab52918, 1:200), TrkA (cs2508, 1:200), p75^{NTR} (cs4201, 1:400) and sortilin (ANT009,1:500). B. Mouse isotype control antibody was used in immunohistochemistry for proNGF (6E10E7, 2.5µg/ml). Scale=25µm

A**B**

Supplementary Figure 1: Negative controls for immunohistochemistry. A. Rabbit isotype control antibody was used in immunohistochemistry for NGF (ab52918, 1:200), TrkA (cs2508, 1:200), p75^{NTR} (cs4201, 1:400) and sortilin (ANT009, 1:500). B. Mouse isotype control antibody was used in immunohistochemistry for proNGF (6E10E7, 2.5 μg/ml). Scale=25 μm