

The RLR/NLR expression and pro-inflammatory activity of tissue mast cells are regulated by cathelicidin LL-37 and defensin hBD-2

Justyna Agier,¹ Sylwia Różalska,² Magdalena Wiktorska,³ Paulina Żelechowska,¹ Joanna Pastwińska,^{1,4} Ewa Brzezińska-Błaszczyk¹

¹Department of Experimental Immunology, Faculty of Health Sciences, Medical University of Lodz, Lodz, Poland

²Department of Industrial Microbiology and Biotechnology, Faculty of Biology and Environmental Protection, University of Lodz, Lodz, Poland

³Department of Molecular Cell Mechanisms, Faculty of Health Sciences, Medical University of Lodz, Lodz, Poland

⁴Laboratory of Cellular Immunology, Institute of Medical Biology, Polish Academy of Sciences, Lodz, Poland

Keywords: Mast cells · NOD1 · NOD2 · RIG-I · Defensin · Cathelicidin

Correspondence:

Prof. Ewa Brzezińska-Błaszczyk, PhD

Department of Experimental Immunology, Medical University of Lodz

Pomorska 251, 92-213 Lodz, Poland

Phone: +48 42 272 57 97

e-mail: ewab@csk.umed.lodz.pl

Supplemental Figure S1

Full length blots of Figure 1b. Red dotted lines show the cropping locations.

