

Multiple mechanisms contributing to ciprofloxacin resistance among Gram negative bacteria causing infections to cancer patients

Samira M. Hamed¹, Walid F. Elkhatib^{2*}, Hadir A. El-Mahallawy³, Mai M. Helmy⁴, Mohamed S. Ashour⁵ and Khaled M. A. Aboshanab²

¹Department of Microbiology and Immunology, Faculty of Pharmacy, October University for Modern Sciences and Arts, 6th of October, Egypt

²Department of Microbiology and Immunology, Faculty of Pharmacy, Ain Shams University, African Union Organization St. Abbassia, Cairo 11566, Egypt

³Department of Clinical Pathology, National Cancer Institute, Cairo University, Cairo, Egypt

⁴Department of Microbiology and Immunology, Faculty of Medicine, Zagazig University, Zagazig, Egypt

⁵Department of Microbiology and Immunology, Faculty of Pharmacy, Al-Azhar University, Cairo, Egypt


*Corresponding author: Prof. Dr. Walid F. Elkhatib

Postal address: ¹Department of Microbiology & Immunology, Faculty of Pharmacy, Ain Shams University, African Union Organization St. Abbassia, Cairo 11566, Egypt

Tel: +202-24051120, Fax: +202-24051107

Email: walid-elkhatib@pharma.asu.edu.eg

Supplementary Figure S1



Supplementary Figure S1. Dendrograms constructed using ERIC-PCR patterns of the *qnr*-positive isolates showing similarity percentages calculated by the dice similarity index. (A) *qnr*-positive *K. pneumoniae* isolates and (B) *qnr*-positive *E. coli* isolates. BAL, bronchoalveolar lavage; CIP MIC, ciprofloxacin MIC (mg/L); PMQR, plasmid mediated quinolone resistance determinants; MDF, MIC decrease factor of ciprofloxacin. Dashed lines are hypothetical lines showing 85% similarity.