

Changing patterns of adult asthma incidence: results from the National Health Insurance Service–National Sample Cohort (NHIS-NSC) database in Korea

Ji-Yeon Shin, MD, PhD¹, Kyoung-Hee Sohn, MD², Ji Eun Shin, PhD³, Mira Park, PhD³,
Jiseun Lim, MD, PhD³, Jin Yong Lee, MD, PhD^{4,*}, Min-Suk Yang, MD, PhD^{2,5,*}

¹Department of Preventive Medicine, School of Medicine, Kyungpook National University, Daegu, Korea,


²Division of Allergy and Clinical Immunology, Department of Internal Medicine, Seoul National University College of Medicine, Seoul, Korea,

³Department of Preventive Medicine, School of Medicine, Eulji University, Daejeon, Korea,

⁴Public Health Medical Service, Boramae Medical Center, Seoul National University College of Medicine, Seoul, Korea,

⁵Department of Internal Medicine, Boramae Medical Center, Seoul National University College of Medicine, Seoul, Korea

Supplementary Figure 1. Trends of age-adjusted incidence rates of asthma in adults (per 1000 person-years) from 2004 to 2012, by sex



Year	p<0.001
Sex	p<0.001
Year*sex	p<0.001

Supplementary Figure 2. Trends in age- and sex-adjusted incidence rates of asthma in adults (per 1000 Person-Years) from 2004 to 2012, by income level (5 groups)

