



Common midwife toad ranaviruses replicate first in the oral cavity of smooth newts (*Lissotriton vulgaris*) and show distinct strain-associated pathogenicity

Bernardo Saucedo^{1,2}, Trenton W.J. Garner³, Natasja Kruithof^{1,2}, Steven J.R. Allain^{4,5}, Mark J. Goodman⁵, Raymond J. Cranfield⁶, Chris Sergeant³, Diego A.Vergara⁷, Marja J.L. Kik^{1,2}, María J. Forzán⁸, Steven J. van Beurden¹, and Andrea Gröne^{1,2\$}

¹Utrecht University, Utrecht, the Netherlands; ²Dutch Wildlife Health Centre, Utrecht, the Netherlands; ³Zoological Society of London, United Kingdom; ⁴Durrell Institute of Ecology and Evolution, University of Kent, UK, ⁵Cambridgeshire and Peterborough Amphibian and Reptile Group, United Kingdom, ⁶Essex Amphibian and Reptile Group, United Kingdom, ⁷National Autonomous University of Mexico, Mexico, ⁸Cornell University, College of Veterinary Medicine USA.



Supplementary Figure 1. Original gel photo from the β actin PCR of intestinal samples from the three experimental groups. Bands of varying intensity were observed in all fields



Supplementary Figure S2. Original gel photo from smooth newt intestinal samples tested for *Ranavirus* infection. Top Row: CMTV-NL I, medium row: CMTV-NL III, bottom row: control group.