

Supplementary Information

Age-Dependent Variation of Lamina Cribrosa Displacement During the Standardized Valsalva Maneuver

Yong Woo Kim, MD^{1,2}, Dong Hyun Lee, MD², Hyung Bin Lim, MD², Baek-Lok Oh, MD^{1,2}, Young Kook Kim, MD¹, Michael J. A. Girard, PhD^{3,4}, Jean Martial Mari, PhD⁵, Ki Ho Park, MD, PhD¹, Jin Wook Jeoung, MD, PhD¹




¹Department of Ophthalmology, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, Korea

²Department of Ophthalmology, Armed Forces Capital Hospital, Seongnam, Korea

³Department of Biomedical Engineering, National University of Singapore, Singapore

⁴Singapore Eye Research Institute, Singapore

⁵University of French Polynesia, Tahiti, French Polynesia



Supplementary Figure S1. Distribution of expiratory pressures during IOP measurement, optic disc cube scan and 21 HD-lines scan between younger and older age groups.

The expiratory pressure during IOP measurement and OCT scans revealed right-skewed distributions. There were no significant differences in expiratory pressures during IOP measurement ($P= 0.61$), optic disc cube scan ($P= 0.89$), or 21 HD-line scan ($P= 0.46$) between younger (groups A to C, $n = 33$) and older age groups (groups D and E, $n = 21$).