

Supplemental Information:

General synovitis score and immunologic synovitis score reflect clinical disease activity in patients with advanced stage rheumatoid arthritis

Tobias Schmidt^{1,5}, Aurélie Najm^{2,3}, Haider Mussawy¹, Rolf Burghardt¹, Nicola Oehler¹, Veit Krenn⁴,
Wolfgang R  ther¹ and Andreas Niemeier¹

¹Department of Orthopaedics, University Medical Center Hamburg-Eppendorf, Martinistra  e 52, 20246
Hamburg, Germany


² Rheumatology Unit, Nantes University Hospital, 44093 Nantes, France.

³ INSERM UMR1238, Nantes University, 44093 Nantes, France

⁴ Department of Pathology, Institute of Pathology, Max-Planck-Stra  e 18, 54296 Trier, Germany


⁵Institute of Osteology and Biomechanics IOBM, University Medical Center Hamburg-Eppendorf,
Martinistra  e 52, 20246 Hamburg, Germany

Supplemental Figure 1:



Supplemental Figure 1: Correlation of different clinical disease activity scores
Disease Activity Score 28 (DAS28), clinical disease activity index (CDAI) and Simplified Disease Activity Index (SDAI) were correlated with each other

Supplemental Figure 2:



Supplemental Figure 2: Patients with refractory monosynovitis display low infiltration of T cells, B cells and neutrophils but high macrophages infiltration

Patients with low clinical disease activity (CDAI ≤ 10) and high synovial membrane inflammation (GSS > 5 and IMSYC > 11) were compared to patients with moderate or high clinical disease activity (CDAI > 10) for (b) GSS and IMSYC and (c) CD 3, CD15, CD20 and (d) CD68 and Ki67 expression. * p values < 0.05 were considered statistically significant.