

SUPPLEMENTARY INFORMATION

Task-dependent functional organizations of the visual ventral stream


Han-Gue Jo^{1,2,*}, Junji Ito³, Barbara Schulte Holthausen¹, Conrad Baumann⁴, Sonja Grün^{3,5}, Ute Habel^{1,2}, Thilo Kellermann^{1,2}

1. Department of Psychiatry, Psychotherapy and Psychosomatics, Medical Faculty, RWTH Aachen University, 52074 Aachen, Germany
2. JARA-Institute Brain Structure Function Relationship (INM-10), Research Center Jülich and RWTH Aachen University, 52074 Aachen, Germany
3. Institute of Neuroscience and Medicine (INM-6) and Institute for Advanced Simulation (IAS-6), Research Center Jülich, 52425 Jülich, Germany
4. Median Klinik Mecklenburg, 19217 Rehna, Germany
5. Theoretical Systems Neurobiology, Faculty I, RWTH Aachen University, 52056 Aachen, Germany

*, Correspondence: hjo@ukaachen.de



Supplementary figure 1. One stimulus image of a task condition is shown. Koala, train, flower, pin, and elephant are embedded in the scene background image. The background image is a photo taken by one of the authors (MS) of Ito et al. [Sci Rep. 2017 7(1):1082. doi: 10.1038/s41598-017-01076-w], and used with permission. The embedded objects were taken from the Microsoft image gallery.



Supplementary figure 2. Group level t-maps from the conjunction of the three task conditions. Significant BOLD signal changes in response to the scene images were thresholded at voxel level uncorrected $p < 0.001$ and cluster-level FEW-corrected $p < 0.05$. The numbers on the bottom indicate z coordinate in MNI space.