

Supplementary information

Association between the relative abundance of gastric microbiota and the risk of gastric cancer: a case-control study

Madhawa Neranjan Gunathilake¹, Jeonghee Lee², Il Ju Choi³, Young-Il Kim³, Yongju Ahn⁴, Chanhyeok Park⁴, Jeongseon Kim^{2*}

¹Department of Cancer Control and Population Health, Graduate School of Cancer Science and Policy, Goyang-si, 10408, Gyeonggi-do, South Korea


²Department of Cancer Biomedical Science, Graduate School of Cancer Science and Policy, Goyang-si, 10408, Gyeonggi-do, South Korea

³Center for Gastric Cancer, National Cancer Center Hospital, National Cancer Center, Goyang-si, 10408, Gyeonggi-do, South Korea

⁴Microbiome Division, Theragen Etex, 145 Gwanggyo-ro, Gyeongtong-gu, Suwon-si, Gyeonggi-do, 16229, South Korea

***Corresponding Author:**

Jeongseon Kim, PhD, Department of Cancer Biomedical Science, Graduate School of Cancer Science and Policy, Goyang-si, 10408, Gyeonggi-do, South Korea. Tel: +82-31-920-2570; Fax: 82-31-920-2579; E-mail: jskim@ncc.re.kr



Supplementary Figure S1. Receiver operating characteristic (ROC) curve for GC prediction using two linear combinations of the four bacterial species. An analysis using two linear combinations of four bacterial species showed 79.7% sensitivity and 67.1% specificity. The AUC was 77.7% indicating that there is a 77.7% chance that the model will be able to distinguish between positive and negative classes. The best cutoff point was chosen as 0.492 in order to find a balance between sensitivity and specificity.