

Loss of the serine protease HTRA1 impairs smooth muscle cells maturation

Ralph Klose,¹ Alexander Prinz,¹ Fabian Tetzlaff,^{1,2} Eva-Maria Weis,¹ Iris Moll,¹ Juan Rodriguez-Vita,¹ Chio Oka,³ Thomas Korff,⁴ Andreas Fischer^{1,2,5,#}

¹Division Vascular Signaling and Cancer (A270), German Cancer Research Center, 69120 Heidelberg, Germany.

²European Center for Angioscience, Medical Faculty Mannheim, Heidelberg University, 68167 Mannheim, Germany.

³Laboratory of Gene Function in Animals, Nara Institute of Science and Technology, 8916-5 Takayama, Ikoma, Nara 630-0192, Japan.

⁴Institute of Physiology and Pathophysiology, Heidelberg University and Deutsches Zentrum für Herz-Kreislauf-Forschung e.V. (DZHK), Partner site Heidelberg/Mannheim, 69120 Heidelberg, Germany.

⁵Medical Clinic I, Endocrinology and Clinical Chemistry, Heidelberg University Hospital, 69120 Heidelberg, Germany.



***Correspondence:**

Andreas Fischer, M.D.
German Cancer Research Center
A270, Im Neuenheimer Feld 280
69120 Heidelberg, Germany
Email: a.fischer@dkfz.de
Phone: +49 6221 42 4150
FAX: +49 6221 42 4159





Running title: HTRA1 is essential for vascular homeostasis

Supplementary Figure 1

a







c





Supplementary Figure 1 (continued)

d





Supplementary Figure 2



a




c



f




Supplementary Figure 3




Supplementary Figure 3 (continued)


f



NOTCH3-ICD



SM22a



GAPDH

