


Optimized Isolation of Extracellular Vesicles From Various Organic Sources Using Aqueous Two Phase System

Oğuz Kaan Kırbaş¹, Batuhan Turhan Bozkurt¹, Ayla Burçin Asutay¹, Beyza Mat¹, Bihter Özdemir¹, Dilek Öztürkoğlu¹, Hülya Ölmez², Zeynep İşlek¹, Fikrettin Şahin¹, and Pakize Neslihan Taşlı^{1*}


¹Yeditepe University, Faculty of Engineering and Architecture, Department of Genetics and Bioengineering, Kayisdagi St., 34755, Istanbul, Turkey.

²Tubitak Marmara Research Center, Baris Dist,41470, Kocaeli, Turkey.


*pneslihan.tasli@yeditepe.edu.tr



Supplementary figure 1: NTA (Nanoparticle Tracking Analysis) data shows concentration (particles/mL) and size (nm) distribution of EVs isolated from Cell Culture, Serum, Parasite Culture and Plant Lysate samples.



Supplementary figure 2: Wide range SEM images of EVs isolated from Cell Culture, Serum, Parasite Culture and Plant Lysate samples.



Supplementary figure 3: Complete gel images for Western Blot Analysis of EVs isolated from Cell Culture, Serum, Parasite Culture and Plant Lysate samples.